

Complexity I

- From past discussion, we know
decidable \rightarrow computationally solvable
- However, this does not mean it is solvable in practice
- The running time may be just too long

Example 1

- $A = \{0^k 1^k \mid k \geq 0\}$

What's the # steps by a 1-tape TM to process a string?

- Remember the procedure
 - 1 check if input is $0^* 1^*$
 - 2 repeat until no 0 or 1 scan, cross off single 0 and 1
 - 3 if 0 or 1 remains, reject otherwise, accept
- How much time?
Need to count # steps

Analysis I

- worst-case analysis
longest time for all inputs
- average-case analysis
- Usually it is easier to do worst-case analysis
- We use a function

$$f : N \rightarrow N$$

to represent the number of steps

N : natural number

n : length of input, $f(n)$: number of steps

Big-O I

- A way to understand the running time of the algorithm when it is run on large inputs
- Consider

$$f(n) = 6n^3 + 5$$

We have

$$n \rightarrow \infty, 6n^3 + 5 \approx 6n^3$$

- $O(f(n)) = O(n^3)$

How about 6 ?

$$6n^3 \text{ vs. } n^3$$

$$6n^3 \text{ vs. } n^4$$

Big-O II

Only things involved with n are important

- Definition:

$$f(n) = O(g(n))$$

if

$$\exists c, n_0, \forall n \geq n_0, f(n) \leq cg(n).$$

Example 1

- Consider

$$f(n) = 6n^3 + 5$$

We have

$$6n^3 + 5 \leq 7n^3 \text{ after } n \geq 2$$

That is, we choose

$$c = 7 \text{ and } n_0 = 2$$

Thus

$$f(n) = O(n^3)$$

Example II

- $f(n) = O(n^4)$ as

$$6n^3 + 5 \leq 7n^4, \text{ after } n \geq 2$$

- But $f(n) \neq O(n^2)$

$$6n^3 + 5 \leq cn^2$$

cannot always hold because we can choose large n such that

$$n^3 > cn^2$$

Example III

- Formally we have the following opposite statement of the definition:

$$\forall c, n_0, \exists n \geq n_0, f(n) > cg(n)$$

Example 7.4 I

- Consider

$$f(n) = 3n \log_2 n + 5n \log_2 \log_2 n$$

We prove

$$f(n) = O(n \log n)$$

by

$$\begin{aligned} \log_2 \log_2 n &\leq \log_2 n \text{ from } \log_2 n \leq n \\ f(n) &\leq 8n \log_2 n = 8n \log_2 b \log_b n \end{aligned}$$

Example 7.4 II

- Note that

$$\frac{\log_2 n}{\log_2 b} = \log_b n$$

- So we write

$$f(n) = O(n \log n)$$

as there is no need to write $\log_2 n$

Other properties I

- We have

$$O(n^2) + O(n) = O(n^2)$$

- Formally,

$$\begin{aligned} f(n) = O(n^2), g(n) = O(n) \\ \Rightarrow f(n) + g(n) = O(n^2) \end{aligned}$$

Other properties II

- Proof

$$\exists c_1, n_1, \forall n \geq n_1, f(n) \leq c_1 n^2$$

$$\exists c_2, n_2, \forall n \geq n_2, g(n) \leq c_2 n$$

Then

$$f(n) + g(n) \leq c_1 n^2 + c_2 n \leq (c_1 + c_2) n^2$$

after $n \geq \max(n_1, n_2)$

Thus we choose

$$c = c_1 + c_2 \text{ and } n_0 = \max(n_1, n_2)$$

Other properties III

- Definition:

$$f(n) = 2^{O(n)}$$

if $\exists c, n_0$ such that

$$f(n) \leq 2^{cn}, \forall n \geq n_0$$

- $O(1)$: $\exists c, n_0$ such that

$$f(n) \leq c1, \forall n \geq n_0$$

Thus

$$f(n) \leq \max\{f(1), \dots, f(n_0 - 1), c\}, \forall n$$

Other properties IV

- That is,

$f(n)$ always \leq a constant