### Homework #0

# **1** Probability and Statistics

(1) (combinatorics)

Let C(N, K) = 1 for K = 0 or K = N, and C(N, K) = C(N - 1, K) + C(N - 1, K - 1) for  $N \ge 1$ . Prove that  $C(N, K) = \frac{N!}{K!(N-K)!}$  for  $N \ge 1$  and  $0 \le K \le N$ .

#### (2) (counting)

What is the probability of getting exactly 4 heads when flipping 10 fair coins?

What is the probability of getting a full house (XXXYY) when randomly drawing 5 cards out of a deck of 52 cards?

(3) (conditional probability)

If your friend flipped a fair coin three times, and tell you that one of the tosses resulted in head, what is the probability that all three tosses resulted in heads?

(4) (Bayes theorem)

A program selects a random integer X like this: a random bit is first generated uniformly. If the bit is 0, X is drawn uniformly from  $\{0, 1, ..., 7\}$ ; otherwise, X is drawn uniformly from  $\{0, -1, -2, -3\}$ . If we get an X from the program with |X| = 1, what is the probability that X is negative?

(5) (union/intersection)

If P(A) = 0.3 and P(B) = 0.4, what is the maximum possible value of  $P(A \cap B)$ ? what is the minimum possible value of  $P(A \cup B)$ ? what is the minimum possible value of  $P(A \cup B)$ ?

# 2 Linear Algebra

(1) (rank)

What is the rank of 
$$\begin{pmatrix} 1 & 2 & 1 \\ 1 & 0 & 3 \\ 1 & 1 & 2 \end{pmatrix}$$
?

(2) (inverse)

What is the inverse of  $\begin{pmatrix} 0 & 2 & 4 \\ 2 & 4 & 2 \\ 3 & 3 & 1 \end{pmatrix}$ ?

(3) (eigenvalues/eigenvectors)

What are the eigenvalues and eigenvectors of  $\begin{pmatrix} 3 & 1 & 1 \\ 2 & 4 & 2 \\ -1 & -1 & 1 \end{pmatrix}$ ?

- (4) (singular value decomposition)
  - (a) For a real matrix M, let  $M = U\Sigma V^T$  be its singular value decomposition. Define  $M^{\dagger} = V\Sigma^{\dagger}U^T$ , where  $\Sigma^{\dagger}[i][j] = \frac{1}{\Sigma[i][j]}$  when  $\Sigma[i][j]$  is nonzero, and 0 otherwise. Prove that  $MM^{\dagger}M = M$ .
  - (b) If M is invertible, prove that  $M^{\dagger} = M^{-1}$ .

(5) (PD/PSD)

A symmetric real matrix A is positive definite (PD) *iff*  $\mathbf{x}^T A \mathbf{x} > 0$  for all  $\mathbf{x} \neq \mathbf{0}$ , and positive semidefinite (PSD) if ">" is changed to " $\geq$ ". Prove:

- (a) For any real matrix Z,  $ZZ^T$  is PSD.
- (b) A symmetric A is PD *iff* all eigenvalues of A are strictly positive.
- (6) (inner product)

Consider  $\mathbf{x} \in \mathbb{R}^d$  and some  $\mathbf{u} \in \mathbb{R}^d$  with  $\|\mathbf{u}\| = 1$ . What is the maximum value of  $\mathbf{u}^T \mathbf{x}$ ? What  $\mathbf{u}$  results in the maximum value? What is the minimum value of  $\mathbf{u}^T \mathbf{x}$ ? What  $\mathbf{u}$  results in the minimum value? What is the minimum value of  $|\mathbf{u}^T \mathbf{x}|$ ? What  $\mathbf{u}$  results in the minimum value?

# 3 Calculus

(1) (differential and partial differential)

Let 
$$f(x) = \ln(1 + e^{-2x})$$
. What is  $\frac{df(x)}{dx}$ ? Let  $g(x, y) = e^x + e^{2y} + e^{3xy^2}$ . What is  $\frac{\partial g(x, y)}{\partial y}$ ?

(2) (chain rule)

Let f(x,y) = xy,  $x(u,v) = \cos(u+v)$ ,  $y(u,v) = \sin(u-v)$ . What is  $\frac{\partial f}{\partial v}$ ?

- (3) (gradient and Hessian) Let  $E(u, v) = (ue^v - 2ve^{-u})^2$ . Calculate the gradient  $\nabla E$  and the Hessian  $\nabla^2 E$  at u = 1 and v = 1.
- (4) (Taylor's expansion) Let  $E(u, v) = (ue^v - 2ve^{-u})^2$ . Write down the second-order Taylor's expansion of E around u = 1and v = 1.
- (5) (optimization) For some given A > 0, B > 0, solve

$$\min_{\alpha} A e^{\alpha} + B e^{-2\alpha}.$$

(6) (vector calculus)

Let **w** be a vector in  $R^d$  and  $E(\mathbf{w}) = \frac{1}{2}\mathbf{w}^T \mathbf{A}\mathbf{w} + \mathbf{b}^T \mathbf{w}$  for some symmetric matrix A and vector **b**. Prove that the gradient  $\nabla E(\mathbf{w}) = \mathbf{A}\mathbf{w} + \mathbf{b}$  and the Hessian  $\nabla^2 E(\mathbf{w}) = \mathbf{A}$ .