Graph Coloring

- k-COLORING asks if the nodes of a graph can be colored with $\leq k$ colors such that no two adjacent nodes have the same color.
- 2-COLORING is in P (why?).
- But 3-COLORING is NP-complete (see next page).
- k-COLORING is NP-complete for $k \geq 3$ (why?).

3-COLORING Is NP-Complete[a]

- We will reduce NAESAT to 3-COLORING.
- We are given a set of clauses $C_1, C_2, \ldots, C_m$ each with 3 literals.
- The boolean variables are $x_1, x_2, \ldots, x_n$.
- We shall construct a graph G such that it can be colored with colors $\{0, 1, 2\}$ if and only if all the clauses can be NAE-satisfied.

[a] Karp (1972).

The Proof (continued)

- Every variable x_i is involved in a triangle $[\, a, x_i, \neg x_i \,]$ with a common node a.
- Each clause $C_i = (c_{i1} \vee c_{i2} \vee c_{i3})$ is also represented by a triangle

$$[\, c_{i1}, c_{i2}, c_{i3} \,].$$

  - Node c_{ij} with the same label as one in some triangle $[\, a, x_k, \neg x_k \,]$ represent *distinct* nodes.
- There is an edge between c_{ij} and the node that represents the jth literal of C_i.

Construction for $\cdots \wedge (x_1 \vee \neg x_2 \vee \neg x_3) \wedge \cdots$



The Proof (continued)

Suppose the graph is 3-colorable.

- Assume without loss of generality that node a takes the color 2.
- A triangle must use up all 3 colors.
- As a result, one of x_i and $\neg x_i$ must take the color 0 and the other 1.

The Proof (continued)

- Treat 1 as `true` and 0 as `false`.[a]
  - We were dealing only with those triangles with the a node, not the clause triangles.
- The resulting truth assignment is clearly contradiction free.
- As each clause triangle contains one color 1 and one color 0, the clauses are NAE-satisfied.

[a] The opposite also works.

The Proof (continued)

Suppose the clauses are NAE-satisfiable.

- Color node a with color 2.
- Color the nodes representing literals by their truth values (color 0 for `false` and color 1 for `true`).
  - We were dealing only with those triangles with the a node, not the clause triangles.

The Proof (concluded)

- For each clause triangle:
  - Pick any two literals with opposite truth values.
  - Color the corresponding nodes with 0 if the literal is `true` and 1 if it is `false`.
  - Color the remaining node with color 2.
- The coloring is legitimate.
  - If literal w of a clause triangle has color 2, then its color will never be an issue.
  - If literal w of a clause triangle has color 1, then it must be connected up to literal w with color 0.
  - If literal w of a clause triangle has color 0, then it must be connected up to literal w with color 1.

TRIPARTITE MATCHING

- We are given three sets B, G, and H, each containing n elements.
- Let $T \subseteq B \times G \times H$ be a ternary relation.
- TRIPARTITE MATCHING asks if there is a set of n triples in T, none of which has a component in common.
  - Each element in B is matched to a different element in G and different element in H.

Theorem 39 (Karp (1972)) TRIPARTITE MATCHING *is NP-complete.*

Related Problems

- We are given a family $F = \{S_1, S_2, \ldots, S_n\}$ of subsets of a finite set U and a budget B.
- SET COVERING asks if there exists a set of B sets in F whose union is U.
- SET PACKING asks if there are B disjoint sets in F.
- Assume $|U| = 3m$ for some $m \in \mathbb{N}$ and $|S_i| = 3$ for all i.
- EXACT COVER BY 3-SETS asks if there are m sets in F that are disjoint and have U as their union.



SET COVERING



SET PACKING

Related Problems (concluded)

Corollary 40 SET COVERING, SET PACKING, *and* EXACT COVER BY 3-SETS *are all NP-complete.*

The KNAPSACK Problem

- There is a set of n items.
- Item i has value $v_i \in \mathbb{Z}^+$ and weight $w_i \in \mathbb{Z}^+$.
- We are given $K \in \mathbb{Z}^+$ and $W \in \mathbb{Z}^+$.
- KNAPSACK asks if there exists a subset $S \subseteq \{1, 2, \ldots, n\}$ such that $\sum_{i \in S} w_i \leq W$ and $\sum_{i \in S} v_i \geq K$.
  - We want to achieve the maximum satisfaction within the budget.

KNAPSACK Is NP-Complete

- KNAPSACK $\in$ NP: Guess an S and verify the constraints.
- We assume $v_i = w_i$ for all i and $K = W$.
- KNAPSACK now asks if a subset of $\{v_1, v_2, \ldots, v_n\}$ adds up to exactly K.
  - Picture yourself as a radio DJ.
  - Or a person trying to control the calories intake.
- We shall reduce EXACT COVER BY 3-SETS to KNAPSACK.

The Proof (continued)

- We are given a family $F = \{S_1, S_2, \ldots, S_n\}$ of size-3 subsets of $U = \{1, 2, \ldots, 3m\}$.
- EXACT COVER BY 3-SETS asks if there are m disjoint sets in F that cover the set U.
- Think of a set as a bit vector in $\{0, 1\}^{3m}$.
  - `001100010` means the set $\{3, 4, 8\}$, and `110010000` means the set $\{1, 2, 5\}$.
- Our goal is $\overbrace{\mathtt{11} \cdots \mathtt{1}}^{3m}$.

The Proof (continued)

- A bit vector can also be considered as a binary *number*.
- Set union resembles addition.
  - $\texttt{001100010} + \texttt{110010000} = \texttt{111110010}$, which denotes the set $\{1, 2, 3, 4, 5, 8\}$, as desired.
- Trouble occurs when there is *carry*.
  - $\texttt{001100010} + \texttt{001110000} = \texttt{010010010}$, which denotes the set $\{2, 5, 8\}$, not the desired $\{3, 4, 5, 8\}$.

The Proof (continued)

- Carry may also lead to a situation where we obtain our solution $\texttt{11}\cdots\texttt{1}$ with more than m sets in F.
  - $\texttt{001100010} + \texttt{001110000} + \texttt{101100000} + \texttt{000001101} = \texttt{111111111}$.
  - But this "solution" $\{1, 3, 4, 5, 6, 7, 8, 9\}$ does not correspond to an exact cover.
  - And it uses 4 sets instead of the required 3.[a]
- To fix this problem, we enlarge the base just enough so that there are no carries.
- Because there are n vectors in total, we change the base from 2 to $n + 1$.

[a] Thanks to a lively class discussion on November 20, 2002.

The Proof (continued)

- Set v_i to be the $(n+1)$-ary number corresponding to the bit vector encoding S_i.
- Now in base $n+1$, if there is a set S such that $\sum_{v_i \in S} v_i = \overbrace{11 \cdots 1}^{3m}$, then every bit position must be contributed by exactly one v_i and $|S| = m$.
- Finally, set

$$K = \sum_{j=0}^{3m-1} (n+1)^j = \overbrace{11 \cdots 1}^{3m} \quad (\text{base } n+1).$$

The Proof (continued)

- Suppose F admits an exact cover, say $\{S_1, S_2, \ldots, S_m\}$.
- Then picking $S = \{v_1, v_2, \ldots, v_m\}$ clearly results in

$$v_1 + v_2 + \cdots + v_m = \overbrace{11 \cdots 1}^{3m}.$$

  - It is important to note that the meaning of addition (+) is independent of the base.[a]
  - It is just regular addition.

[a] Contributed by Mr. Kuan-Yu Chen (R92922047) on November 3, 2004.

The Proof (concluded)

- On the other hand, suppose there exists an S such that $\sum_{v_i \in S} v_i = \overbrace{11 \cdots 1}^{3m}$ in base $n+1$.
- The no-carry property implies that $|S| = m$ and $\{S_i : v_i \in S\}$ is an exact cover.

An Example

- Let $m = 3$, $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$, and

$$\begin{aligned} S_1 &= \{1, 3, 4\}, \\ S_2 &= \{2, 3, 4\}, \\ S_3 &= \{2, 5, 6\}, \\ S_4 &= \{6, 7, 8\}, \\ S_5 &= \{7, 8, 9\}. \end{aligned}$$

- Note that $n = 5$, as there are 5 S_i's.

An Example (concluded)

- Our reduction produces

$$
\begin{aligned}
K &= \sum_{j=0}^{3\times 3-1} 6^j = \overbrace{11\cdots 1}^{3\times 3} \quad \text{(base 6)}, \\
v_1 &= \texttt{101100000}, \\
v_2 &= \texttt{011100000}, \\
v_3 &= \texttt{010011000}, \\
v_4 &= \texttt{000001110}, \\
v_5 &= \texttt{000000111}.
\end{aligned}
$$

- Note $v_1 + v_3 + v_5 = K$.
- Indeed, $S_1 \cup S_3 \cup S_5 = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$, an exact cover by 3-sets.

BIN PACKINGS

- We are given N positive integers $a_1, a_2, \ldots, a_N$, an integer C (the capacity), and an integer B (the number of bins).
- BIN PACKING asks if these numbers can be partitioned into B subsets, each of which has total sum at most C.
- Think of packing bags at the check-out counter.

Theorem 41 BIN PACKING *is NP-complete.*

INTEGER PROGRAMMING

- INTEGER PROGRAMMING asks whether a system of linear inequalities with integer coefficients has an integer solution.
  - LINEAR PROGRAMMING asks whether a system of linear inequalities with integer coefficients has a *rational* solution.

INTEGER PROGRAMMING Is NP-Complete[a]

- SET COVERING can be expressed by the inequalities $Ax \geq \vec{1}$, $\sum_{i=1}^{n} x_i \leq B$, $0 \leq x_i \leq 1$, where
  - x_i is one if and only if S_i is in the cover.
  - A is the matrix whose columns are the bit vectors of the sets $S_1, S_2, \ldots$.
  - $\vec{1}$ is the vector of 1s.
- This shows INTEGER PROGRAMMING is NP-hard.
- Many NP-complete problems can be expressed as an INTEGER PROGRAMMING problem.

[a] Papadimitriou (1981).

Easier or Harder?[a]

- Adding restrictions on the allowable *problem instances* will not make a problem harder.
  - We are now solving a subset of problem instances.
  - The INDEPENDENT SET proof (p. 277) and the KNAPSACK proof (p. 322).
  - SAT to 2SAT (easier by p. 264).
  - CIRCUIT VALUE to MONOTONE CIRCUIT VALUE (equally hard by p. 241).

[a]Thanks to a lively class discussion on October 29, 2003.

Easier or Harder? (concluded)

- Adding restrictions on the allowable *solutions* may make a problem easier, as hard, or harder.
- It is problem dependent.
  - MIN CUT to BISECTION WIDTH (harder by p. 303).
  - LINEAR PROGRAMMING to INTEGER PROGRAMMING (harder by p. 332).
  - SAT to NAESAT (equally hard by p. 272) and MAX CUT to MAX BISECTION (equally hard by p. 301).
  - 3-COLORING to 2-COLORING (easier by p. 309).

coNP and Function Problems

coNP

- By definition, coNP is the class of problems whose complement is in NP.
- NP is the class of problems that have succinct certificates (recall Proposition 30 on p. 251).
- coNP is therefore the class of problems that have succinct disqualifications:
  - A "no" instance of a problem in coNP possesses a short proof of its being a "no" instance.
  - Only "no" instances have such proofs.

coNP (continued)

- Suppose L is a coNP problem.
- There exists a polynomial-time nondeterministic algorithm M such that:
  - If $x \in L$, then $M(x) =$ "yes" for all computation paths.
  - If $x \notin L$, then $M(x) =$ "no" for some computation path.


$x \in L$
yes
yes
yes
yes
yes


$x \notin L$
no
yes
yes
no
yes

coNP (concluded)

- Clearly $\mathrm{P} \subseteq \mathrm{coNP}$.
- It is not known if

$$\mathrm{P} = \mathrm{NP} \cap \mathrm{coNP}.$$

  - Contrast this with

$$\mathrm{R} = \mathrm{RE} \cap \mathrm{coRE}$$

    (see Proposition 11 on p. 124).

Some coNP Problems

- VALIDITY $\in$ coNP.
  - If ϕ is not valid, it can be disqualified very succinctly: a truth assignment that does not satisfy it.
- SAT COMPLEMENT $\in$ coNP.
  - The disqualification is a truth assignment that satisfies it.
- HAMILTONIAN PATH COMPLEMENT $\in$ coNP.
  - The disqualification is a Hamiltonian path.
- OPTIMAL TSP (D) $\in$ coNP.[a]
  - The disqualification is a tour with a length $< B$.

[a] Asked by Mr. Che-Wei Chang (R95922093) on September 27, 2006.

An Alternative Characterization of coNP

Proposition 42 *Let $L \subseteq \Sigma^*$ be a language. Then $L \in coNP$ if and only if there is a polynomially decidable and polynomially balanced relation R such that*

$$L = \{x : \forall y \, (x, y) \in R\}.$$

(As on p. 250, we assume $|\, y \,| \leq |\, x \,|^k$ for some k.)

- $\bar{L} = \{x : (x, y) \in \neg R \text{ for some } y\}$.
- Because $\neg R$ remains polynomially balanced, $\bar{L} \in \text{NP}$ by Proposition 30 (p. 251).
- Hence $L \in \text{coNP}$ by definition.

coNP Completeness

Proposition 43 *L is NP-complete if and only if its complement $\bar{L} = \Sigma^* - L$ is coNP-complete.*

Proof ($\Rightarrow$; the $\Leftarrow$ part is symmetric)

- Let $\bar{L}'$ be any coNP language.
- Hence $L' \in$ NP.
- Let R be the reduction from L' to L.
- So $x \in L'$ if and only if $R(x) \in L$.
- So $x \in \bar{L}'$ if and only if $R(x) \in \bar{L}$.
- R is a reduction from $\bar{L}'$ to $\bar{L}$.

Some coNP-Complete Problems

- SAT COMPLEMENT is coNP-complete.
  - SAT COMPLEMENT is the complement of SAT.
- VALIDITY is coNP-complete.
  - ϕ is valid if and only if $\neg\phi$ is not satisfiable.
  - The reduction from SAT COMPLEMENT to VALIDITY is hence easy.
- HAMILTONIAN PATH COMPLEMENT is coNP-complete.

Possible Relations between P, NP, coNP

1. $\mathrm{P} = \mathrm{NP} = \mathrm{coNP}$.
2. $\mathrm{NP} = \mathrm{coNP}$ but $\mathrm{P} \neq \mathrm{NP}$.
3. $\mathrm{NP} \neq \mathrm{coNP}$ and $\mathrm{P} \neq \mathrm{NP}$.
   - This is current "consensus."

coNP Hardness and NP Hardness[a]

Proposition 44 *If a coNP-hard problem is in NP, then NP = coNP.*

- Let $L \in \text{NP}$ be coNP-hard.
- Let NTM M decide L.
- For any $L' \in \text{coNP}$, there is a reduction R from L' to L.
- $L' \in \text{NP}$ as it is decided by NTM $M(R(x))$.
  - Alternatively, NP is closed under complement.
- Hence $\text{coNP} \subseteq \text{NP}$.
- The other direction $\text{NP} \subseteq \text{coNP}$ is symmetric.

[a] Brassard (1979); Selman (1978).

coNP Hardness and NP Hardness (concluded)

Similarly,

Proposition 45 *If an NP-hard problem is in coNP, then* $NP = coNP$.

Hence NP-complete problems are unlikely to be in coNP and coNP-complete problems are unlikely to be in NP.

The Primality Problem

- An integer p is **prime** if $p > 1$ and all positive numbers other than 1 and p itself cannot divide it.
- PRIMES asks if an integer N is a prime number.
- Dividing N by $2, 3, \ldots, \sqrt{N}$ is *not* efficient.
  - The length of N is only $\log N$, but $\sqrt{N} = 2^{0.5 \log N}$.
- A polynomial-time algorithm for PRIMES was not found until 2002 by Agrawal, Kayal, and Saxena!
- We will focus on efficient "probabilistic" algorithms for PRIMES (used in *Mathematica*, e.g.).

```
if n = a^b for some a, b > 1 then
   return "composite";
end if
for r = 2, 3, ..., n − 1 do
   if gcd(n, r) > 1 then
      return "composite";
   end if
   if r is a prime then
      Let q be the largest prime factor of r − 1;
      if q ≥ 4√r log n and n^((r−1)/q) ≠ 1 mod r then
         break; {Exit the for-loop.}
      end if
   end if
end for{r − 1 has a prime factor q ≥ 4√r log n.}
for a = 1, 2, ..., 2√r log n do
   if (x − a)^n ≠ (x^n − a) mod (x^r − 1) in Z_n[x] then
      return "composite";
   end if
end for
return "prime"; {The only place with "prime" output.}
```

DP

- DP $\equiv$ NP $\cap$ coNP is the class of problems that have succinct certificates and succinct disqualifications.
  - Each "yes" instance has a succinct certificate.
  - Each "no" instance has a succinct disqualification.
  - No instances have both.
- P $\subseteq$ DP.
- We will see that PRIMES $\in$ DP.
  - In fact, PRIMES $\in$ P as mentioned earlier.

Primitive Roots in Finite Fields

Theorem 46 (Lucas and Lehmer (1927)) [a] *A number* $p > 1$ *is prime if and only if there is a number* $1 < r < p$ *(called the* **primitive root** *or* **generator***) such that*

1. $r^{p-1} = 1 \bmod p$, *and*
2. $r^{(p-1)/q} \neq 1 \bmod p$ *for all prime divisors* q *of* $p - 1$.

- We will prove the theorem later.

[a] François Edouard Anatole Lucas (1842–1891); Derrick Henry Lehmer (1905–1991).

Pratt's Theorem

Theorem 47 (Pratt (1975)) PRIMES $\in NP \cap coNP$.

- PRIMES is in coNP because a succinct disqualification is a divisor.
- Suppose p is a prime.
- p's certificate includes the r in Theorem 46 (p. 351).
- Use recursive doubling to check if $r^{p-1} = 1 \bmod p$ in time polynomial in the length of the input, $\log_2 p$.
- We also need all *prime* divisors of $p - 1$: $q_1, q_2, \ldots, q_k$.
- Checking $r^{(p-1)/q_i} \neq 1 \bmod p$ is also easy.

The Proof (concluded)

- Checking $q_1, q_2, \ldots, q_k$ are all the divisors of $p-1$ is easy.
- We still need certificates for the primality of the q_i's.
- The complete certificate is recursive and tree-like:

$$C(p) = (r; q_1, C(q_1), q_2, C(q_2), \ldots, q_k, C(q_k)).$$

- $C(p)$ can also be checked in polynomial time.
- We next prove that $C(p)$ is succinct.

The Succinctness of the Certificate

Lemma 48 *The length of $C(p)$ is at most quadratic at $5\log_2^2 p$.*

- This claim holds when $p = 2$ or $p = 3$.
- In general, $p - 1$ has $k < \log_2 p$ prime divisors $q_1 = 2, q_2, \ldots, q_k$.
- $C(p)$ requires: 2 parentheses and $2k < 2\log_2 p$ separators (length at most $2\log_2 p$ long), r (length at most $\log_2 p$), $q_1 = 2$ and its certificate 1 (length at most 5 bits), the q_i's (length at most $2\log_2 p$), and the $C(q_i)$s.

The Proof (concluded)

- $C(p)$ is succinct because

$$
\begin{aligned}
|C(p)| &\leq 5\log_2 p + 5 + 5\sum_{i=2}^{k} \log_2^2 q_i \\
&\leq 5\log_2 p + 5 + 5\left(\sum_{i=2}^{k} \log_2 q_i\right)^2 \\
&\leq 5\log_2 p + 5 + 5\log_2^2 \frac{p-1}{2} \\
&< 5\log_2 p + 5 + 5(\log_2 p - 1)^2 \\
&= 5\log_2^2 p + 10 - 5\log_2 p \leq 5\log_2^2 p
\end{aligned}
$$

for $p \geq 4$.

Basic Modular Arithmetics[a]

- Let $m, n \in \mathbb{Z}^+$.
- $m|n$ means m divides n and m is n's **divisor**.
- We call the numbers $0, 1, \ldots, n-1$ the **residue** modulo n.
- The **greatest common divisor** of m and n is denoted $\gcd(m, n)$.
- The r in Theorem 46 (p. 351) is a primitive root of p.
- We now prove the existence of primitive roots and then Theorem 46.

[a] Carl Friedrich Gauss.

Euler's[a] Totient or Phi Function

- Let

$$\Phi(n) = \{m : 1 \leq m < n, \gcd(m, n) = 1\}$$

  be the set of all positive integers less than n that are prime to n (Z_n^* is a more popular notation).
  - $\Phi(12) = \{1, 5, 7, 11\}$.
- Define **Euler's function** of n to be $\phi(n) = |\Phi(n)|$.
- $\phi(p) = p - 1$ for prime p, and $\phi(1) = 1$ by convention.
- Euler's function is not expected to be easy to compute without knowing n's factorization.

[a] Leonhard Euler (1707–1783).


ϕ(n)
500
400
300
200
100
200
300
400
500
n

Two Properties of Euler's Function

The inclusion-exclusion principle[a] can be used to prove the following.

Lemma 49 $\phi(n) = n \prod_{p|n} (1 - \frac{1}{p})$.

- If $n = p_1^{e_1} p_2^{e_2} \cdots p_t^{e_t}$ is the prime factorization of n, then

$$\phi(n) = n \prod_{i=1}^{t} \left(1 - \frac{1}{p_i}\right).$$

Corollary 50 $\phi(mn) = \phi(m)\,\phi(n)$ *if* $\gcd(m, n) = 1$.

[a] See my *Discrete Mathematics* lecture notes.

A Key Lemma

Lemma 51 $\sum_{m|n} \phi(m) = n$.

- Let $\prod_{i=1}^{\ell} p_i^{k_i}$ be the prime factorization of n and consider

$$\prod_{i=1}^{\ell} [\, \phi(1) + \phi(p_i) + \cdots + \phi(p_i^{k_i}) \,]. \tag{4}$$

- Equation (4) equals n because $\phi(p_i^k) = p_i^k - p_i^{k-1}$ by Lemma 49.
- Expand Eq. (4) to yield $\sum_{k_1' \le k_1, \ldots, k_\ell' \le k_\ell} \prod_{i=1}^{\ell} \phi(p_i^{k_i'})$.

The Proof (concluded)

- By Corollary 50 (p. 359),

$$\prod_{i=1}^{\ell} \phi(p_i^{k_i'}) = \phi\left(\prod_{i=1}^{\ell} p_i^{k_i'}\right).$$

- Each $\prod_{i=1}^{\ell} p_i^{k_i'}$ is a unique divisor of $n = \prod_{i=1}^{\ell} p_i^{k_i}$.
- Equation (4) becomes

$$\sum_{m|n} \phi(m).$$

The Density Attack for PRIMES



- It works, but does it work well?

Factorization and Euler's Function

- The ratio of numbers $\leq n$ relatively prime to n is $\phi(n)/n$.
- When $n = pq$, where p and q are distinct primes,

$$\frac{\phi(n)}{n} = \frac{pq - p - q + 1}{pq} > 1 - \frac{1}{q} - \frac{1}{p}.$$

  - The "density attack" to factor $n = pq$ hence takes $\Omega(\sqrt{n})$ steps on average when $p \sim q = O(\sqrt{n}\,)$.
  - This running time is exponential: $\Omega(2^{0.5 \log_2 n})$.

The Chinese Remainder Theorem

- Let $n = n_1 n_2 \cdots n_k$, where n_i are pairwise relatively prime.
- For any integers $a_1, a_2, \ldots, a_k$, the set of simultaneous equations

$$\begin{aligned} x &= a_1 \bmod n_1, \\ x &= a_2 \bmod n_2, \\ &\vdots \\ x &= a_k \bmod n_k, \end{aligned}$$

has a unique solution modulo n for the unknown x.

Fermat's "Little" Theorem[a]

Lemma 52 *For all* $0 < a < p$, $a^{p-1} = 1 \bmod p$.

- Consider $a\Phi(p) = \{am \bmod p : m \in \Phi(p)\}$.
- $a\Phi(p) = \Phi(p)$.
  - $a\Phi(p) \subseteq \Phi(p)$ as a remainder must be between 0 and $p - 1$.
  - Suppose $am = am' \bmod p$ for $m > m'$, where $m, m' \in \Phi(p)$.
  - That means $a(m - m') = 0 \bmod p$, and p divides a or $m - m'$, which is impossible.

[a] Pierre de Fermat (1601–1665).

The Proof (concluded)

- Multiply all the numbers in $\Phi(p)$ to yield $(p-1)!$.
- Multiply all the numbers in $a\Phi(p)$ to yield $a^{p-1}(p-1)!$.
- As $a\Phi(p) = \Phi(p)$, $(p-1)! = a^{p-1}(p-1)! \bmod p$.
- Finally, $a^{p-1} = 1 \bmod p$ because $p \not| (p-1)!$.

The Fermat-Euler Theorem[a]

Corollary 53 *For all* $a \in \Phi(n)$, $a^{\phi(n)} = 1 \bmod n$.

- The proof is similar to that of Lemma 52 (p. 365).
- Consider $a\Phi(n) = \{am \bmod n : m \in \Phi(n)\}$.
- $a\Phi(n) = \Phi(n)$.
  - $a\Phi(n) \subseteq \Phi(n)$ as a remainder must be between 0 and $n-1$ and relatively prime to n.
  - Suppose $am = am' \bmod n$ for $m' < m < n$, where $m, m' \in \Phi(n)$.
  - That means $a(m - m') = 0 \bmod n$, and n divides a or $m - m'$, which is impossible.

[a] Proof by Mr. Wei-Cheng Cheng (R93922108) on November 24, 2004.

The Proof (concluded)

- Multiply all the numbers in $\Phi(n)$ to yield $\prod_{m \in \Phi(n)} m$.
- Multiply all the numbers in $a\Phi(n)$ to yield $a^{\Phi(n)} \prod_{m \in \Phi(n)} m$.
- As $a\Phi(n) = \Phi(n)$,

$$\prod_{m \in \Phi(n)} m = a^{\Phi(n)} \left(\prod_{m \in \Phi(n)} m \right) \bmod n.$$

- Finally, $a^{\Phi(n)} = 1 \bmod n$ because $n \not| \prod_{m \in \Phi(n)} m$.

An Example

- As $12 = 2^2 \times 3$,

$$\phi(12) = 12 \times \left(1 - \frac{1}{2}\right)\left(1 - \frac{1}{3}\right) = 4$$

- In fact, $\Phi(12) = \{1, 5, 7, 11\}$.
- For example,

$$5^4 = 625 = 1 \bmod 12.$$