Primality Tests

- PRIMES asks if a number N is a prime.
- The classic algorithm tests if $k \mid N$ for $k = 2, 3, \ldots, \sqrt{N}$.
- But it runs in $\Omega(2^{n/2})$ steps, where $n = |N| = \log_2 N$.

The Density Attack for PRIMES

```
Pick k ∈ {2, ..., N − 1} randomly; {Assume N > 2.}
if k | N then
    return "N is composite";
else
    return "N is a prime";
end if
```

Analysis[a]

- Suppose $N = PQ$, a product of 2 primes.
- The probability of success is

$$< 1 - \frac{\phi(N)}{N} = 1 - \frac{(P-1)(Q-1)}{PQ} = \frac{P+Q-1}{PQ}.$$

- In the case where $P \approx Q$, this probability becomes

$$< \frac{1}{P} + \frac{1}{Q} \approx \frac{2}{\sqrt{N}}.$$

- This probability is exponentially small.

[a] See also p. 363.

The Fermat Test for Primality

Fermat's "little" theorem on p. 365 suggests the following primality test for any given number p:

```
Pick a number a randomly from {1, 2, ..., N − 1};
if a^(N−1) ≠ 1 mod N then
    return "N is composite";
else
    return "N is probably a prime";
end if
```

The Fermat Test for Primality (concluded)

- Unfortunately, there are composite numbers called **Carmichael numbers** that will pass the Fermat test for *all* $a \in \{1, 2, \ldots, N-1\}$.
- There are infinitely many Carmichael numbers.[a]

[a] Alford, Granville, and Pomerance (1992).

Square Roots Modulo a Prime

- Equation $x^2 = a \bmod p$ has at most two (distinct) roots by Lemma 54 (p. 370).
  - The roots are called **square roots**.
  - Numbers a with square roots and $\gcd(a, p) = 1$ are called **quadratic residues**.
    - They are $1^2 \bmod p, 2^2 \bmod p, \ldots, (p-1)^2 \bmod p$.
- We shall show that a number either has two roots or has none, and testing which one is true is trivial.
- There are no known efficient *deterministic* algorithms to find the roots.

Euler's Test

Lemma 60 (Euler) *Let p be an odd prime and $a \neq 0 \bmod p$.*

1. *If $a^{(p-1)/2} = 1 \bmod p$, then $x^2 = a \bmod p$ has two roots.*
2. *If $a^{(p-1)/2} \neq 1 \bmod p$, then $a^{(p-1)/2} = -1 \bmod p$ and $x^2 = a \bmod p$ has no roots.*

- Let r be a primitive root of p.
- By Fermat's "little" theorem, $r^{(p-1)/2}$ is a square root of 1, so $r^{(p-1)/2} = \pm 1 \bmod p$.
- But as r is a primitive root, $r^{(p-1)/2} \neq 1 \bmod p$.
- Hence $r^{(p-1)/2} = -1 \bmod p$.

The Proof (continued)

- Suppose $a = r^{2j}$ for some $1 \leq j \leq (p-1)/2$.
- Then $a^{(p-1)/2} = r^{j(p-1)} = 1 \bmod p$ and its two *distinct* roots are $r^j, -r^j (= r^{j+(p-1)/2})$.
  - If $r^j = -r^j \bmod p$, then $2r^j = 0 \bmod p$, which implies $r^j = 0 \bmod p$, a contradiction.
- As $1 \leq j \leq (p-1)/2$, there are $(p-1)/2$ such a's.

The Proof (continued)

- Each such a has 2 distinct square roots.
- The square roots of all the a's are distinct.
  - The square roots of different a's must be different.
- Hence the set of *square roots* is $\{1, 2, \ldots, p-1\}$.
  - Because there are $(p-1)/2$ such a's and each a has two square roots.
- As a result, $a = r^{2j}$, $1 \leq j \leq (p-1)/2$, are all the quadratic residues.

The Proof (concluded)

- If $a = r^{2j+1}$, then it has no roots because all the square roots have been taken.
- Now,

$$a^{(p-1)/2} = [\, r^{(p-1)/2} \,]^{2j+1} = (-1)^{2j+1} = -1 \bmod p.$$

The Legendre Symbol[a] and Quadratic Residuacity Test

- By Lemma 60 (p. 426) $a^{(p-1)/2} \bmod p = \pm 1$ for $a \neq 0 \bmod p$.
- For odd prime p, define the **Legendre symbol** $(a \mid p)$ as

$$(a \mid p) = \begin{cases} 0 & \text{if } p \mid a, \\ 1 & \text{if } a \text{ is a quadratic residue modulo } p, \\ -1 & \text{if } a \text{ is a } \textbf{quadratic nonresidue} \text{ modulo } p. \end{cases}$$

- Euler's test implies $a^{(p-1)/2} = (a \mid p) \bmod p$ for any odd prime p and any integer a.
- Note that $(ab|p) = (a|p)(b|p)$.

[a] Andrien-Marie Legendre (1752–1833).

Gauss's Lemma

Lemma 61 (Gauss) *Let p and q be two odd primes. Then $(q|p) = (-1)^m$, where m is the number of residues in $R = \{iq \bmod p : 1 \leq i \leq (p-1)/2\}$ that are greater than $(p-1)/2$.*

- All residues in R are distinct.
  - If $iq = jq \bmod p$, then $p|(j-i)\,q$ or $p|q$.
- No two elements of R add up to p.
  - If $iq + jq = 0 \bmod p$, then $p|(i+j)$ or $p|q$.
  - But neither is possible.

The Proof (continued)

- Consider the set R' of residues that result from R if we replace each of the m elements $a \in R$ such that $a > (p-1)/2$ by $p - a$.
  - This is equivalent to performing $-a \bmod p$.
- All residues in R' are now at most $(p-1)/2$.
- In fact, $R' = \{1, 2, \ldots, (p-1)/2\}$ (see illustration next page).
  - Otherwise, two elements of R would add up to p, which has been shown to be impossible.



$p = 7$ and $q = 5$.

The Proof (concluded)

- Alternatively, $R' = \{\pm iq \bmod p : 1 \leq i \leq (p-1)/2\}$, where exactly m of the elements have the minus sign.
- Take the product of all elements in the two representations of R'.
- So $[(p-1)/2]! = (-1)^m q^{(p-1)/2}[(p-1)/2]! \bmod p$.
- Because $\gcd([(p-1)/2]!, p) = 1$, the above implies

$$1 = (-1)^m q^{(p-1)/2} \bmod p.$$

Legendre's Law of Quadratic Reciprocity[a]

- Let p and q be two odd primes.
- The next result says their Legendre symbols are distinct if and only if both numbers are 3 mod 4.

Lemma 62 (Legendre (1785), Gauss)

$$(p|q)(q|p) = (-1)^{\frac{p-1}{2}\frac{q-1}{2}}.$$

[a]First stated by Euler in 1751. Legendre (1785) did not give a correct proof. Gauss proved the theorem when he was 19. He gave at least 6 different proofs during his life. The 152nd proof appeared in 1963.

The Proof (continued)

- Sum the elements of R' in the previous proof in mod2.
- On one hand, this is just $\sum_{i=1}^{(p-1)/2} i \bmod 2$.
- On the other hand, the sum equals

$$\begin{aligned} & \sum_{i=1}^{(p-1)/2} \left(qi - p \left\lfloor \frac{iq}{p} \right\rfloor \right) + mp \bmod 2 \\ = & \left(q \sum_{i=1}^{(p-1)/2} i - p \sum_{i=1}^{(p-1)/2} \left\lfloor \frac{iq}{p} \right\rfloor \right) + mp \bmod 2. \end{aligned}$$

  - Signs are irrelevant under mod2.
  - m is as in Lemma 61 (p. 431).

The Proof (continued)

- Ignore odd multipliers to make the sum equal

$$\left(\sum_{i=1}^{(p-1)/2} i - \sum_{i=1}^{(p-1)/2} \left\lfloor \frac{iq}{p} \right\rfloor \right) + m \bmod 2.$$

- Equate the above with $\sum_{i=1}^{(p-1)/2} i \bmod 2$ to obtain

$$m = \sum_{i=1}^{(p-1)/2} \left\lfloor \frac{iq}{p} \right\rfloor \bmod 2.$$

The Proof (concluded)

- $\sum_{i=1}^{(p-1)/2} \lfloor \frac{iq}{p} \rfloor$ is the number of integral points under the line $y = (q/p)\,x$ for $1 \le x \le (p-1)/2$.
- Gauss's lemma (p. 431) says $(q|p) = (-1)^m$.
- Repeat the proof with p and q reversed.
- So $(p|q) = (-1)^{m'}$, where m' is the number of integral points *above* the line $y = (q/p)\,x$ for $1 \le y \le (q-1)/2$.
- As a result, $(p|q)(q|p) = (-1)^{m+m'}$.
- But $m + m'$ is the total number of integral points in the $\frac{p-1}{2} \times \frac{q-1}{2}$ rectangle, which is $\frac{p-1}{2}\,\frac{q-1}{2}$.

Eisenstein's Rectangle



$p = 11$ and $q = 7$.

The Jacobi Symbol[a]

- The Legendre symbol only works for odd *prime* moduli.
- The **Jacobi symbol** $(a \mid m)$ extends it to cases where m is not prime.
- Let $m = p_1 p_2 \cdots p_k$ be the prime factorization of m.
- When $m > 1$ is odd and $\gcd(a, m) = 1$, then

$$(a|m) = \prod_{i=1}^{k} (a \mid p_i).$$

- Define $(a \mid 1) = 1$.

[a] Carl Jacobi (1804–1851).

Properties of the Jacobi Symbol

The Jacobi symbol has the following properties, for arguments for which it is defined.

1. $(ab \mid m) = (a \mid m)(b \mid m)$.
2. $(a \mid m_1 m_2) = (a \mid m_1)(a \mid m_2)$.
3. If $a = b \bmod m$, then $(a \mid m) = (b \mid m)$.
4. $(-1 \mid m) = (-1)^{(m-1)/2}$ (by Lemma 61 on p. 431).
5. $(2 \mid m) = (-1)^{(m^2-1)/8}$ (by Lemma 61 on p. 431).
6. If a and m are both odd, then $(a \mid m)(m \mid a) = (-1)^{(a-1)(m-1)/4}$.

Calculation of $(2200|999)$

Similar to the Euclidean algorithm and does *not* require factorization.

$$
\begin{aligned}
& (202|999) = (-1)^{(999^2-1)/8}(101|999) \\
= \quad & (-1)^{124750}(101|999) = (101|999) \\
= \quad & (-1)^{(100)(998)/4}(999|101) = (-1)^{24950}(999|101) \\
= \quad & (999|101) = (90|101) = (-1)^{(101^2-1)/8}(45|101) \\
= \quad & (-1)^{1275}(45|101) = -(45|101) \\
= \quad & -(-1)^{(44)(100)/4}(101|45) = -(101|45) = -(11|45) \\
= \quad & -(-1)^{(10)(44)/4}(45|11) = -(45|11) \\
= \quad & -(1|11) = -(11|1) = -1.
\end{aligned}
$$

A Result Generalizing Proposition 10.3 in the Textbook

Theorem 63 *The group of set* $\Phi(n)$ *under multiplication* mod n *has a primitive root if and only if* n *is either 1, 2, 4,* p^k*, or* $2p^k$ *for some nonnegative integer* k *and and odd prime* p*.*

This result is essential in the proof of the next lemma.

The Jacobi Symbol and Primality Test[a]

Lemma 64 *If* $(M|N) = M^{(N-1)/2} \bmod N$ *for all* $M \in \Phi(N)$*, then* N *is prime. (Assume* N *is odd.)*

- Assume $N = mp$, where p is an odd prime, $\gcd(m, p) = 1$, and $m > 1$ (not necessarily prime).
- Let $r \in \Phi(p)$ such that $(r \,|\, p) = -1$.
- The Chinese remainder theorem says that there is an $M \in \Phi(N)$ such that

$$\begin{aligned} M &= r \bmod p, \\ M &= 1 \bmod m. \end{aligned}$$

[a]Mr. Clement Hsiao (R88526067) pointed out that the textbook's proof in Lemma 11.8 is incorrect while he was a senior in January 1999.

The Proof (continued)

- By the hypothesis,

$$M^{(N-1)/2} = (M \mid N) = (M \mid p)(M \mid m) = -1 \bmod N.$$

- Hence

$$M^{(N-1)/2} = -1 \bmod m.$$

- But because $M = 1 \bmod m$,

$$M^{(N-1)/2} = 1 \bmod m,$$

  a contradiction.

The Proof (continued)

- Second, assume that $N = p^a$, where p is an odd prime and $a \geq 2$.
- By Theorem 63 (p. 443), there exists a primitive root r modulo p^a.
- From the assumption,

$$M^{N-1} = \left[M^{(N-1)/2} \right]^2 = (M|N)^2 = 1 \bmod N$$

for all $M \in \Phi(N)$.

The Proof (continued)

- As $r \in \Phi(N)$ (prove it), we have

$$r^{N-1} = 1 \bmod N.$$

- As r's exponent modulo $N = p^a$ is $\phi(N) = p^{a-1}(p-1)$,

$$p^{a-1}(p-1) \mid N-1,$$

  which implies that $p \mid N-1$.

- But this is impossible given that $p \mid N$.

The Proof (continued)

- Third, assume that $N = mp^a$, where p is an odd prime, $\gcd(m, p) = 1$, $m > 1$ (not necessarily prime), and a is even.
- The proof mimics that of the second case.
- By Theorem 63 (p. 443), there exists a primitive root r modulo p^a.
- From the assumption,

$$M^{N-1} = \left[M^{(N-1)/2} \right]^2 = (M|N)^2 = 1 \bmod N$$

for all $M \in \Phi(N)$.

The Proof (continued)

- In particular,

$$M^{N-1} = 1 \bmod p^a \tag{6}$$

  for all $M \in \Phi(N)$.

- The Chinese remainder theorem says that there is an $M \in \Phi(N)$ such that

$$\begin{aligned} M &= r \bmod p^a, \\ M &= 1 \bmod m. \end{aligned}$$

- Because $M = r \bmod p^a$ and Eq. (6),

$$r^{N-1} = 1 \bmod p^a.$$

The Proof (concluded)

- As r's exponent modulo $N = p^a$ is $\phi(N) = p^{a-1}(p-1)$,

$$p^{a-1}(p-1) \mid N-1,$$

  which implies that $p \mid N-1$.

- But this is impossible given that $p \mid N$.

The Number of Witnesses to Compositeness

Theorem 65 (Solovay and Strassen (1977)) *If N is an odd composite, then $(M|N) \neq M^{(N-1)/2} \bmod N$ for at least half of $M \in \Phi(N)$.*

- By Lemma 64 (p. 444) there is at least one $a \in \Phi(N)$ such that $(a|N) \neq a^{(N-1)/2} \bmod N$.
- Let $B = \{b_1, b_2, \ldots, b_k\} \subseteq \Phi(N)$ be the set of *all* distinct residues such that $(b_i|N) = b_i^{(N-1)/2} \bmod N$.
- Let $aB = \{ab_i \bmod N : i = 1, 2, \ldots, k\}$.

The Proof (concluded)

- $|aB| = k$.
  - $ab_i = ab_j \bmod N$ implies $N|a(b_i - b_j)$, which is impossible because $\gcd(a, N) = 1$ and $N > |b_i - b_j|$.
- $aB \cap B = \emptyset$ because

$$(ab_i)^{(N-1)/2} = a^{(N-1)/2} b_i^{(N-1)/2} \neq (a|N)(b_i|N) = (ab_i|N).$$

- Combining the above two results, we know

$$\frac{|\,B\,|}{\phi(N)} \leq 0.5.$$

```
if N is even but N ≠ 2 then
    return "N is composite";
else if N = 2 then
    return "N is a prime";
end if
Pick M ∈ {2, 3, ..., N − 1} randomly;
if gcd(M, N) > 1 then
    return "N is a composite";
else
    if (M|N) ≠ M^((N−1)/2) mod N then
        return "N is composite";
    else
        return "N is a prime";
    end if
end if
```

Analysis

- The algorithm certainly runs in polynomial time.
- There are no false positives (for COMPOSITENESS).
  - When the algorithm says the number is composite, it is always correct.
- The probability of a false negative is at most one half.
  - When the algorithm says the number is a prime, it may err.
  - If the input is composite, then the probability that the algorithm errs is one half.
- The error probability can be reduced but not eliminated.

The Improved Density Attack for COMPOSITENESS



Randomized Complexity Classes; RP

- Let N be a polynomial-time precise NTM that runs in time $p(n)$ and has 2 nondeterministic choices at each step.
- N is a **polynomial Monte Carlo Turing machine** for a language L if the following conditions hold:
  - If $x \in L$, then at least half of the $2^{p(n)}$ computation paths of N on x halt with "yes" where $n = |\, x \,|$.
  - If $x \notin L$, then all computation paths halt with "no."
- The class of all languages with polynomial Monte Carlo TMs is denoted **RP** (**randomized polynomial time**).[a]

[a] Adleman and Manders (1977).

Comments on RP

- Nondeterministic steps can be seen as fair coin flips.
- There are no false positive answers.
- The probability of false negatives, $1 - \epsilon$, is at most 0.5.
- But any constant between 0 and 1 can replace 0.5.
  - By repeating the algorithm $k = \lceil -\frac{1}{\log_2 1-\epsilon} \rceil$ times, the probability of false negatives becomes $(1 - \epsilon)^k \leq 0.5$.
- In fact, ϵ can be arbitrarily close to 0 as long as it is of the order $1/p(n)$ for some polynomial $p(n)$.
  - $-\frac{1}{\log_2 1-\epsilon} = O(\frac{1}{\epsilon}) = O(p(n))$.

Where RP Fits

- $\mathrm{P} \subseteq \mathrm{RP} \subseteq \mathrm{NP}$.
  - A deterministic TM is like a Monte Carlo TM except that all the coin flips are ignored.
  - A Monte Carlo TM is an NTM with extra demands on the number of accepting paths.
- COMPOSITENESS $\in$ RP; PRIMES $\in$ coRP; PRIMES $\in$ RP.[a]
  - In fact, PRIMES $\in$ P.[b]
- $\mathrm{RP} \cup \mathrm{coRP}$ is another "plausible" notion of efficient computation.

[a] Adleman and Huang (1987).
[b] Agrawal, Kayal, and Saxena (2002).

ZPP[a] (Zero Probabilistic Polynomial)

- The class **ZPP** is defined as RP $\cap$ coRP.
- A language in ZPP has *two* Monte Carlo algorithms, one with no false positives and the other with no false negatives.
- If we repeatedly run both Monte Carlo algorithms, *eventually* one definite answer will come (unlike RP).
  - A *positive* answer from the one without false positives.
  - A *negative* answer from the one without false negatives.

[a] Gill (1977).

The ZPP Algorithm (**Las Vegas**)

```
{Suppose L ∈ ZPP.}
{N_1 has no false positives, and N_2 has no false negatives.}
while true do
    if N_1(x) = "yes" then
        return "yes";
    end if
    if N_2(x) = "no" then
        return "no";
    end if
end while
```

ZPP (concluded)

- The *expected* running time for the correct answer to emerge is polynomial.
  - The probability that a run of the 2 algorithms does not generate a definite answer is 0.5.
  - Let $p(n)$ be the running time of each run.
  - The expected running time for a definite answer is

$$\sum_{i=1}^{\infty} 0.5^i i p(n) = 2p(n).$$

- Essentially, ZPP is the class of problems that can be solved without errors in expected polynomial time.

Et Tu, RP?

```
{Suppose L ∈ RP.}
{N decides L without false positives.}
while true do
   if N(x) = "yes" then
      return "yes";
   end if
   {But what to do here?}
end while
```

- You eventually get a "yes" if $x \in L$.
- But how to get a "no" when $x \notin L$?
- You have to sacrifice either correctness or bounded running time.

Large Deviations

- Suppose you have a *biased* coin.
- One side has probability $0.5 + \epsilon$ to appear and the other $0.5 - \epsilon$, for some $0 < \epsilon < 0.5$.
- But you do not know which is which.
- How to decide which side is the more likely—with high confidence?
- Answer: Flip the coin many times and pick the side that appeared the most times.
- Question: Can you quantify the confidence?

The Chernoff Bound[a]

Theorem 66 (Chernoff (1952)) *Suppose $x_1, x_2, \ldots, x_n$ are independent random variables taking the values 1 and 0 with probabilities p and $1-p$, respectively. Let $X = \sum_{i=1}^{n} x_i$. Then for all $0 \leq \theta \leq 1$,*

$$\text{prob}[\, X \geq (1+\theta)\, pn \,] \leq e^{-\theta^2 pn/3}.$$

- The probability that the deviate of a **binomial random variable** from its expected value $E[\, X \,] = E[\, \sum_{i=1}^{n} x_i \,] = pn$ decreases exponentially with the deviation.
- The Chernoff bound is asymptotically optimal.

[a] Herman Chernoff (1923–).

The Proof

- Let t be any positive real number.
- Then

$$\text{prob}[\, X \geq (1+\theta)\, pn \,] = \text{prob}[\, e^{tX} \geq e^{t(1+\theta)\, pn} \,].$$

- Markov's inequality (p. 405) generalized to real-valued random variables says that

$$\text{prob}\left[\, e^{tX} \geq kE[\, e^{tX} \,] \,\right] \leq 1/k.$$

- With $k = e^{t(1+\theta)\, pn} / E[\, e^{tX} \,]$, we have

$$\text{prob}[\, X \geq (1+\theta)\, pn \,] \leq e^{-t(1+\theta)\, pn} E[\, e^{tX} \,].$$

The Proof (continued)

- Because $X = \sum_{i=1}^{n} x_i$ and x_i's are independent,

$$E[\, e^{tX} \,] = (E[\, e^{tx_1} \,])^n = [\, 1 + p(e^t - 1) \,]^n.$$

- Substituting, we obtain

$$\begin{aligned} \text{prob}[\, X \geq (1+\theta)\, pn \,] &\leq e^{-t(1+\theta)\, pn} [\, 1 + p(e^t - 1) \,]^n \\ &\leq e^{-t(1+\theta)\, pn} e^{pn(e^t - 1)} \end{aligned}$$

as $(1+a)^n \leq e^{an}$ for all $a > 0$.

The Proof (concluded)

- With the choice of $t = \ln(1+\theta)$, the above becomes

$$\text{prob}[\, X \geq (1+\theta)\, pn \,] \leq e^{pn[\,\theta - (1+\theta)\ln(1+\theta)\,]}.$$

- The exponent expands to $-\frac{\theta^2}{2} + \frac{\theta^3}{6} - \frac{\theta^4}{12} + \cdots$ for $0 \leq \theta \leq 1$, which is less than

$$-\frac{\theta^2}{2} + \frac{\theta^3}{6} \leq \theta^2 \left(-\frac{1}{2} + \frac{\theta}{6} \right) \leq \theta^2 \left(-\frac{1}{2} + \frac{1}{6} \right) = -\frac{\theta^2}{3}.$$

Power of the Majority Rule

From prob$[\, X \leq (1-\theta)\, pn \,] \leq e^{-\frac{\theta^2}{2} pn}$ (prove it):

Corollary 67 *If* $p = (1/2) + \epsilon$ *for some* $0 \leq \epsilon \leq 1/2$, *then*

$$\text{prob}\left[\sum_{i=1}^{n} x_i \leq n/2\right] \leq e^{-\epsilon^2 n/2}.$$

- The textbook's corollary to Lemma 11.9 seems incorrect.
- Our original problem (p. 463) hence demands $\approx 1.4k/\epsilon^2$ independent coin flips to guarantee making an error with probability at most 2^{-k} with the majority rule.

BPP[a] (Bounded Probabilistic Polynomial)

- The class **BPP** contains all languages for which there is a precise polynomial-time NTM N such that:
  - If $x \in L$, then at least 3/4 of the computation paths of N on x lead to "yes."
  - If $x \notin L$, then at least 3/4 of the computation paths of N on x lead to "no."
- N accepts or rejects by a *clear* majority.

[a] Gill (1977).

Magic 3/4?

- The number 3/4 bounds the probability of a right answer away from 1/2.
- Any constant *strictly* between 1/2 and 1 can be used without affecting the class BPP.
- In fact, 0.5 plus any inverse polynomial between 1/2 and 1,

$$0.5 + \frac{1}{p(n)},$$

can be used.

The Majority Vote Algorithm

Suppose L is decided by N by majority $(1/2) + \epsilon$.

```
for i = 1, 2, ..., 2k + 1 do
    Run N on input x;
end for
if "yes" is the majority answer then
    "yes";
else
    "no";
end if
```

Analysis

- The running time remains polynomial, being $2k + 1$ times N's running time.
- By Corollary 67 (p. 468), the probability of a false answer is at most $e^{-\epsilon^2 k}$.
- By taking $k = \lceil 2/\epsilon^2 \rceil$, the error probability is at most $1/4$.
- As with the RP case, ϵ can be any inverse polynomial, because k remains polynomial in n.

Probability Amplification for BPP

- Let m be the number of random bits used by a BPP algorithm.
  - By definition, m is polynomial in n.
- With $k = \Theta(\log m)$ in the majority vote algorithm, we can lower the error probability to $\leq (3m)^{-1}$.

Aspects of BPP

- BPP is the most comprehensive yet plausible notion of efficient computation.
  - If a problem is in BPP, we take it to mean that the problem can be solved efficiently.
  - In this aspect, BPP has effectively replaced P.
- $(\mathrm{RP} \cup \mathrm{coRP}) \subseteq (\mathrm{NP} \cup \mathrm{coNP})$.
- $(\mathrm{RP} \cup \mathrm{coRP}) \subseteq \mathrm{BPP}$.
- Whether $\mathrm{BPP} \subseteq (\mathrm{NP} \cup \mathrm{coNP})$ is unknown.
- But it is unlikely that $\mathrm{NP} \subseteq \mathrm{BPP}$ (p. 489).

coBPP

- The definition of BPP is symmetric: acceptance by clear majority and rejection by clear majority.
- An algorithm for $L \in \text{BPP}$ becomes one for $\bar{L}$ by reversing the answer.
- So $\bar{L} \in \text{BPP}$ and $\text{BPP} \subseteq \text{coBPP}$.
- Similarly $\text{coBPP} \subseteq \text{BPP}$.
- Hence $\text{BPP} = \text{coBPP}$.
- This approach does not work for RP.
- It did not work for NP either.

BPP and coBPP



"The Good, the Bad, and the Ugly"



Circuit Complexity

- Circuit complexity is based on boolean circuits instead of Turing machines.
- A boolean circuit with n inputs computes a boolean function of n variables.
- By identify `true` with 1 and `false` with 0, a boolean circuit with n inputs accepts certain strings in $\{ 0, 1 \}^n$.
- To relate circuits with arbitrary languages, we need one circuit for each possible input length n.

Formal Definitions

- The **size** of a circuit is the number of *gates* in it.
- A **family of circuits** is an infinite sequence $\mathcal{C} = (C_0, C_1, \ldots)$ of boolean circuits, where C_n has n boolean inputs.
- $L \subseteq \{0,1\}^*$ has **polynomial circuits** if there is a family of circuits $\mathcal{C}$ such that:
  - The size of C_n is at most $p(n)$ for some fixed polynomial p.
  - For input $x \in \{0,1\}^*$, $C_{|x|}$ outputs 1 if and only if $x \in L$.
    - C_n accepts $L \cap \{0,1\}^n$.

Exponential Circuits Contain All Languages

- Theorem 14 (p. 153) implies that there are languages that cannot be solved by circuits of size $2^n/(2n)$.
- But exponential circuits can solve all problems.

Proposition 68 *All decision problems (decidable or otherwise) can be solved by a circuit of size* 2^{n+2}.

- We will show that for any language $L \subseteq \{0,1\}^*$, $L \cap \{0,1\}^n$ can be decided by a circuit of size 2^{n+2}.

The Proof (concluded)

- Define boolean function $f : \{0,1\}^n \rightarrow \{0,1\}$, where

$$f(x_1 x_2 \cdots x_n) = \begin{cases} 1 & x_1 x_2 \cdots x_n \in L, \\ 0 & x_1 x_2 \cdots x_n \notin L. \end{cases}$$

- $f(x_1 x_2 \cdots x_n) = (x_1 \wedge f(1 x_2 \cdots x_n)) \vee (\neg x_1 \wedge f(0 x_2 \cdots x_n))$.
- The circuit size $s(n)$ for $f(x_1 x_2 \cdots x_n)$ hence satisfies

$$s(n) = 4 + 2s(n-1)$$

  with $s(1) = 1$.
- Solve it to obtain $s(n) = 5 \times 2^{n-1} - 4 \leq 2^{n+2}$.

Comments

- Proposition 68 (p. 480) does not contradict anything we knew so far about computation theory.
- Yes, there are only a finite number of circuits with size 2^{n+2}.
- Yes, there are only 2^n possible inputs of length n.
- Yes, those circuits can solve all problems of length n.
- But is there an algorithm to tell which circuit is the correct one?