

Undirected Graphs

- An **undirected graph** $G = (V, E)$ has a finite set of nodes, V , and a set of *undirected* edges, E .
- It is like a directed graph except that the edges have no directions and there are no self-loops.
- Use $[i, j]$ to denote the fact that there is an edge between node i and node j .

Independent Sets

- Let $G = (V, E)$ be an undirected graph.
- $I \subseteq V$.
- I is **independent** if there is no edge between any two nodes $i, j \in I$.
- The INDEPENDENT SET problem: Given an undirected graph and a goal K , is there an independent set of size K ?
- Many applications.


INDEPENDENT SET Is NP-Complete

- This problem is in NP: Guess a set of nodes and verify that it is independent and meets the count.
- We will reduce 3SAT to INDEPENDENT SET.
- If a graph contains a triangle, any independent set can contain at most one node of the triangle.
- The results of the reduction will be graphs whose nodes can be partitioned into m disjoint triangles.

The Proof (continued)

- Let ϕ be an instance of 3SAT with m clauses.
- We will construct graph G with $K = m$.
- Furthermore, ϕ is satisfiable if and only if G has an independent set of size K .
 - Here is the reduction:
 - There is a triangle for each clause with the literals as the nodes.
 - Add edges between x and $\neg x$ for every variable x .

$$(x_1 \vee x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_3) \wedge (\neg x_1 \vee x_2 \vee x_3)$$



Same literal labels that appear in different clauses yield *distinct* nodes.

The Proof (continued)

- Suppose G has an independent set I of size $K = m$.
 - An independent set can contain at most m nodes, one from each triangle.
 - So I contains exactly one node from each triangle.
 - Truth assignment T assigns true to those literals in I .
 - T is consistent because contradictory literals are connected by an edge; hence both cannot be in I .
 - T satisfies ϕ because it has a node from every triangle, thus satisfying every clause.^a

^aThe variables without a truth value can be assigned arbitrarily. Contributed by Mr. Chun-Yuan Chen (R99922119) on November 2, 2010.

The Proof (concluded)

- Suppose a satisfying truth assignment T exists for ϕ .
 - Collect one node from each triangle whose literal is true under T .
 - The choice is arbitrary if there is more than one true literal.
 - This set of m nodes must be independent by construction.
 - * Both literals x and $\neg x$ cannot be assigned true.

Other INDEPENDENT SET-Related NP-Complete Problems

Corollary 37 INDEPENDENT SET *is NP-complete for 4-degree graphs.*

Theorem 38 INDEPENDENT SET *is NP-complete for planar graphs.*

Theorem 39 (Garey and Johnson (1977))
INDEPENDENT SET *is NP-complete for 3-degree planar graphs.*


NODE COVER

- We are given an undirected graph G and a goal K .
- NODE COVER: Is there a set C with K or fewer nodes such that each edge of G has at least one of its endpoints (i.e., incident nodes) in C ?
- Many applications.

NODE COVER Is NP-Complete

Corollary 40 (Karp (1972)) NODE COVER *is NP-complete.*

- I is an independent set of $G = (V, E)$ if and only if $V - I$ is a node cover of G .



CLIQUE

- We are given an undirected graph G and a goal K .
- CLIQUE asks if there is a set C with K nodes such that there is an edge between any two nodes $i, j \in C$.
- Many applications.

Remarks^a

- Are INDEPENDENT SET and NODE COVER NP-complete if K is a constant?
 - No, because one can do an exhaustive search on all the possible node covers or independent sets (both $\binom{n}{k}$ of them, a polynomial).^b
- Are INDEPENDENT SET and NODE COVER NP-complete if K is a linear function of n ?
 - INDEPENDENT SET with $K = n/3$ and NODE COVER with $K = 2n/3$ remain NP-complete by our reductions.


^aContributed by Mr. Ching-Hua Yu (D00921025) on October 30, 2012.

^b $n = |V|$.

CLIQUE Is NP-Complete

Corollary 41 (Karp (1972)) *CLIQUE is NP-complete.*

- Let \bar{G} be the **complement** of G , where $[x, y] \in \bar{G}$ if and only if $[x, y] \notin G$.
- I is a clique in $G \Leftrightarrow I$ is an independent set in \bar{G} .




MIN CUT and MAX CUT

- A **cut** in an undirected graph $G = (V, E)$ is a partition of the nodes into two nonempty sets S and $V - S$.
- The size of a cut $(S, V - S)$ is the number of edges between S and $V - S$.
- MIN CUT $\in P$ by the maxflow algorithm.^a
- MAX CUT asks if there is a cut of size at least K .
 - K is part of the input.

^aIn time $O(|V| \cdot |E|)$ by Orlin (2012).

A Cut of Size 4



MIN CUT and MAX CUT (concluded)

- MAX CUT has applications in circuit layout.
 - The minimum area of a VLSI layout of a graph is not less than the square of its maximum cut size.^a

^aRaspaud, Sýkora, and Vrřo (1995); Mak and Wong (2000).


MAX CUT Is NP-Complete^a

- We will reduce NAESAT to MAX CUT.
- Given an instance ϕ of 3SAT with m clauses, we shall construct a graph $G = (V, E)$ and a goal K .
- Furthermore, there is a cut of size at least K if and only if ϕ is NAE-satisfiable.
- Our graph will have multiple edges between two nodes.
 - Each such edge contributes one to the cut if its nodes are separated.

^aKarp (1972) and Garey, Johnson, and Stockmeyer (1976).


The Proof

- Suppose ϕ 's m clauses are C_1, C_2, \dots, C_m .
- The boolean variables are x_1, x_2, \dots, x_n .
- G has $2n$ nodes: $x_1, x_2, \dots, x_n, \neg x_1, \neg x_2, \dots, \neg x_n$.
- Each clause with 3 distinct literals makes a triangle in G .
- For each clause with two identical literals, there are two parallel edges between the two distinct literals.
- No need to consider clauses with one literal (why?).
- For each variable x_i , add n_i copies of edge $[x_i, \neg x_i]$, where n_i is the number of occurrences of x_i and $\neg x_i$ in ϕ .



The Proof (continued)

- Set $K = 5m$.
- Suppose there is a cut $(S, V - S)$ of size $5m$ or more.
- A clause (a triangle or two parallel edges) contributes at most 2 to a cut no matter how you split it.
- Suppose both x_i and $\neg x_i$ are on the same side of the cut.
- They *together* contribute at most $2n_i$ edges to the cut.
 - They appear in at most n_i different clauses.
 - A clause contributes at most 2 to a cut.



The Proof (continued)


- Either x_i or $\neg x_i$ contributes at most n_i to the cut by the pigeonhole principle.
- Changing the side of that literal does *not decrease* the size of the cut.
- Hence we assume variables are separated from their negations.
- The total number of edges in the cut that join opposite literals x_i and $\neg x_i$ is $\sum_{i=1}^n n_i$.
- But $\sum_{i=1}^n n_i = 3m$ as it is simply the total number of literals.

The Proof (concluded)

- The *remaining* $K - 3m \geq 2m$ edges in the cut must come from the m triangles or parallel edges that correspond to the clauses.
- Each can contribute at most 2 to the cut.^a
- So all are split.
- A split clause means at least one of its literals is true and at least one false.
- The other direction is left as an exercise.


^aSo $K = 5m$.

A Cut That Does Not Meet the Goal $K = 5 \times 3 = 15$



- $(x_1 \vee x_2 \vee x_2) \wedge (x_1 \vee \neg x_3 \vee \neg x_3) \wedge (\neg x_1 \vee \neg x_2 \vee x_3)$.
- The cut size is $13 < 15$.

A Cut That Meets the Goal $K = 5 \times 3 = 15$



- $(x_1 \vee x_2 \vee x_2) \wedge (x_1 \vee \neg x_3 \vee \neg x_3) \wedge (\neg x_1 \vee \neg x_2 \vee x_3)$.
- The cut size is now 15.

Remarks

- We had proved that MAX CUT is NP-complete for multigraphs.
- How about proving the same thing for simple graphs?^a
- How to modify the proof to reduce 4SAT to MAX CUT?^b
- All NP-complete problems are mutually reducible by definition.^c
 - So they are equally hard in this sense.^d

^aContributed by Mr. Tai-Dai Chou (J93922005) on June 2, 2005.

^bContributed by Mr. Chien-Lin Chen (J94922015) on June 8, 2006.

^cContributed by Mr. Ren-Shuo Liu (D98922016) on October 27, 2009.

^dContributed by Mr. Ren-Shuo Liu (D98922016) on October 27, 2009.

MAX BISECTION


- MAX CUT becomes MAX BISECTION if we require that $|S| = |V - S|$.
- It has many applications, especially in VLSI layout.

MAX BISECTION Is NP-Complete

- We shall reduce the more general MAX CUT to MAX BISECTION.
- Add $|V| = n$ **isolated nodes** to G to yield G' .
- G' has $2n$ nodes.
- G' 's goal K is identical to G 's
 - As the new nodes have no edges, they contribute nothing to the cut.
- This completes the reduction.

The Proof (concluded)

- Every cut $(S, V - S)$ of $G = (V, E)$ can be made into a bisection by appropriately allocating the new nodes between S and $V - S$.
- Hence each cut of G can be made a cut of G' of the same size, and vice versa.



BISECTION WIDTH

- BISECTION WIDTH is like MAX BISECTION except that it asks if there is a bisection of size *at most* K (sort of MIN BISECTION).
- Unlike MIN CUT, BISECTION WIDTH is NP-complete.
- We reduce MAX BISECTION to BISECTION WIDTH.
- Given a graph $G = (V, E)$, where $|V|$ is even, we generate the complement of G .
- Given a goal of K , we generate a goal of $n^2 - K$.

The Proof (concluded)

- To show the reduction works, simply notice the following easily verifiable claims.
 - A graph $G = (V, E)$, where $|V| = 2n$, has a bisection of size K if and only if the complement of G has a bisection of size $n^2 - K$.
 - So G has a bisection of size $\geq K$ if and only if its complement has a bisection of size $\leq n^2 - K$.

HAMILTONIAN PATH Is NP-Complete^a

Theorem 42 *Given an undirected graph, the question whether it has a Hamiltonian path is NP-complete.*

^aKarp (1972).

A Hamiltonian Path at IKEA, Covina, California?




TSP (D) Is NP-Complete

Corollary 43 TSP (D) *is NP-complete.*

- Consider a graph G with n nodes.
- Create a weighted complete graph G' with the same nodes as from G follows.
- Set $d_{ij} = 1$ on G' if $[i, j] \in G$ and $d_{ij} = 2$ on G' if $[i, j] \notin G$.
 - Note that G' is a complete graph.
- Set the budget $B = n + 1$.
- This completes the reduction.

TSP (D) Is NP-Complete (continued)

- Suppose G' has a tour of distance at most $n + 1$.
- Then that tour on G' must contain at most one edge with weight 2.
- If a tour on G' contains 1 edge with weight 2, remove that edge to arrive at a Hamiltonian path for G .
- If, on the other hand, a tour on G' contains no edge with weight 2.
- Remove any edge to arrive at a Hamiltonian path for G .



TSP (D) Is NP-Complete (concluded)

- On the other hand, suppose G has Hamiltonian paths.
- Then there is a tour on G' containing at most one edge with weight 2.
- The total cost is then at most $(n - 1) + 2 = n + 1 = B$.
- We conclude that there is a tour of length B or less on G' if and only if G has a Hamiltonian path.

Graph Coloring

- k -COLORING: Can the nodes of a graph be colored with $\leq k$ colors such that no two adjacent nodes have the same color?^a
- 2-COLORING is in P (why?).
- But 3-COLORING is NP-complete (see next page).
- k -COLORING is NP-complete for $k \geq 3$ (why?).
- EXACT- k -COLORING asks if the nodes of a graph can be colored using exactly k colors.
- It remains NP-complete for $k \geq 3$ (why?).

^a k is *not* part of the input; k is part of the problem statement.

3-COLORING Is NP-Complete^a

- We will reduce NAESAT to 3-COLORING.
- We are given a set of clauses C_1, C_2, \dots, C_m each with 3 literals.
- The boolean variables are x_1, x_2, \dots, x_n .
- We shall construct a graph G such that it can be colored with colors $\{0, 1, 2\}$ if and only if all the clauses can be NAE-satisfied.

^aKarp (1972).

The Proof (continued)


- Every variable x_i is involved in a triangle $[a, x_i, \neg x_i]$ with a common node a .
- Each clause $C_i = (c_{i1} \vee c_{i2} \vee c_{i3})$ is also represented by a triangle

$$[c_{i1}, c_{i2}, c_{i3}].$$

- Node c_{ij} with the same label as one in some triangle $[a, x_k, \neg x_k]$ represent *distinct* nodes.
- There is an edge between c_{ij} and the node that represents the j th literal of C_i .
 - Alternative proof: there is an edge between $\neg c_{ij}$ and the node that represents the j th literal of C_i .^a

^aContributed by Mr. Ren-Shuo Liu (D98922016) on October 27, 2009.

Construction for $\dots \wedge (x_1 \vee \neg x_2 \vee \neg x_3) \wedge \dots$



The Proof (continued)

Suppose the graph is 3-colorable.

- Assume without loss of generality that node a takes the color 2.
- A triangle must use up all 3 colors.
- As a result, one of x_i and $\neg x_i$ must take the color 0 and the other 1.

The Proof (continued)

- Treat 1 as true and 0 as false.^a
 - We were dealing only with those triangles with the “*a*” node, not the clause triangles.
- The resulting truth assignment is clearly contradiction free.
- As each clause triangle contains one color 1 and one color 0, the clauses are NAE-satisfied.

^aThe opposite also works.

The Proof (continued)

Suppose the clauses are NAE-satisfiable.

- Color node a with color 2.
- Color the nodes representing literals by their truth values (color 0 for **false** and color 1 for **true**).
 - We were dealing only with those triangles with the “ a ” node, not the clause triangles.

The Proof (continued)

- For each clause triangle:
 - Pick any two literals with opposite truth values.
 - Color the corresponding nodes with 0 if the literal is true and 1 if it is false.
 - Color the remaining node with color 2.

The Proof (concluded)

- The coloring is legitimate.
 - If literal w of a clause triangle has color 2, then its color will never be an issue.
 - If literal w of a clause triangle has color 1, then it must be connected up to literal w with color 0.
 - If literal w of a clause triangle has color 0, then it must be connected up to literal w with color 1.

Algorithms for 3-COLORING and the Chromatic Number $\chi(G)$

- Assume G is 3-colorable.
- There is an algorithm to find a 3-coloring in time $O(3^{n/3}) = 1.4422^n$.^a
- It has been improved to $O(1.3289^n)$.^b

^aLawler (1976).

^bBeigel and Eppstein (2000).

Algorithms for 3-COLORING and the Chromatic Number $\chi(G)$ (concluded)

- The **chromatic number** $\chi(G)$ is the smallest number of colors needed to color a graph G .
- There is an algorithm to find $\chi(G)$ in time $O((4/3)^{n/3}) = 2.4422^n$.^a
- It can be improved to $O((4/3 + 3^{4/3}/4)^n) = O(2.4150^n)$ ^b and $2^n n^{O(1)}$.^c
- Computing $\chi(G)$ cannot be easier than 3-COLORING.^d

^aLawler (1976).

^bEppstein (2003).

^cKoivisto (2006).

^dContributed by Mr. Ching-Hua Yu (D00921025) on October 30, 2012.

TRIPARTITE MATCHING

- We are given three sets B , G , and H , each containing n elements.
- Let $T \subseteq B \times G \times H$ be a ternary relation.
- TRIPARTITE MATCHING asks if there is a set of n triples in T , none of which has a component in common.
 - Each element in B is matched to a different element in G and different element in H .

Theorem 44 (Karp (1972)) TRIPARTITE MATCHING *is NP-complete.*