

## Brownian Bridge Approach to Pricing Barrier Options

- We desire an unbiased estimate which can be calculated efficiently.
- The above-mentioned payoff should be multiplied by the probability  $p$  that a continuous sample path does *not* hit the barrier conditional on the sampled prices.
- This methodology is called the Brownian bridge approach.
- Formally, we have

$$p \equiv \text{Prob}[S(t) < H, 0 \leq t \leq T \mid S(t_0), S(t_1), \dots, S(t_n)].$$

## Brownian Bridge Approach to Pricing Barrier Options (continued)

- As a barrier is hit over a time interval if and only if the maximum stock price over that period is at least  $H$ ,

$$p = \text{Prob} \left[ \max_{0 \leq t \leq T} S(t) < H \mid S(t_0), S(t_1), \dots, S(t_n) \right].$$

- Luckily, the conditional distribution of the maximum over a time interval given the beginning and ending stock prices is known.

## Brownian Bridge Approach to Pricing Barrier Options (continued)

**Lemma 22** Assume  $S$  follows  $dS/S = \mu dt + \sigma dW$  and define

$$\zeta(x) \equiv \exp \left[ -\frac{2 \ln(x/S(t)) \ln(x/S(t + \Delta t))}{\sigma^2 \Delta t} \right].$$

(1) If  $H > \max(S(t), S(t + \Delta t))$ , then

$$\text{Prob} \left[ \max_{t \leq u \leq t + \Delta t} S(u) < H \mid S(t), S(t + \Delta t) \right] = 1 - \zeta(H).$$

(2) If  $h < \min(S(t), S(t + \Delta t))$ , then

$$\text{Prob} \left[ \min_{t \leq u \leq t + \Delta t} S(u) > h \mid S(t), S(t + \Delta t) \right] = 1 - \zeta(h).$$

## Brownian Bridge Approach to Pricing Barrier Options (continued)

- Lemma 22 gives the probability that the barrier is not hit in a time interval, given the starting and ending stock prices.
- For our up-and-out call, choose  $n = 1$ .
- As a result,

$$p = \begin{cases} 1 - \exp \left[ -\frac{2 \ln(H/S(0)) \ln(H/S(T))}{\sigma^2 T} \right], & \text{if } H > \max(S(0), S(T)), \\ 0, & \text{otherwise.} \end{cases}$$

## Brownian Bridge Approach to Pricing Barrier Options (continued)

```
1:  $C := 0$ ;  
2: for  $i = 1, 2, 3, \dots, N$  do  
3:    $P := S \times e^{(r-q-\sigma^2/2)T + \sigma\sqrt{T} \xi()};$   
4:   if  $(S < H$  and  $P < H)$  or  $(S > H$  and  $P > H)$  then  
5:      $C := C + \max(P - X, 0) \times \left\{ 1 - \exp \left[ -\frac{2 \ln(H/S) \times \ln(H/P)}{\sigma^2 T} \right] \right\};$   
6:   end if  
7: end for  
8: return  $C e^{-rT} / N$ ;
```

## Brownian Bridge Approach to Pricing Barrier Options (concluded)

- The idea can be generalized.
- For example, we can handle more complex barrier options.
- Consider an up-and-out call with barrier  $H_i$  for the time interval  $(t_i, t_{i+1}]$ ,  $0 \leq i < n$ .
- This option thus contains  $n$  barriers.
- Multiply the probabilities for the  $n$  time intervals to obtain the desired probability adjustment term.

## Variance Reduction

- The statistical efficiency of Monte Carlo simulation can be measured by the variance of its output.
- If this variance can be lowered without changing the expected value, fewer replications are needed.
- Methods that improve efficiency in this manner are called variance-reduction techniques.
- Such techniques become practical when the added costs are outweighed by the reduction in sampling.

## Variance Reduction: Antithetic Variates

- We are interested in estimating  $E[g(X_1, X_2, \dots, X_n)]$ .
- Let  $Y_1$  and  $Y_2$  be random variables with the same distribution as  $g(X_1, X_2, \dots, X_n)$ .

- Then

$$\text{Var} \left[ \frac{Y_1 + Y_2}{2} \right] = \frac{\text{Var}[Y_1]}{2} + \frac{\text{Cov}[Y_1, Y_2]}{2}.$$

- $\text{Var}[Y_1]/2$  is the variance of the Monte Carlo method with two independent replications.
- The variance  $\text{Var}[(Y_1 + Y_2)/2]$  is smaller than  $\text{Var}[Y_1]/2$  when  $Y_1$  and  $Y_2$  are negatively correlated.



## Variance Reduction: Antithetic Variates (continued)

- For each simulated sample path  $X$ , a second one is obtained by *reusing* the random numbers on which the first path is based.
- This yields a second sample path  $Y$ .
- Two estimates are then obtained: One based on  $X$  and the other on  $Y$ .
- If  $N$  independent sample paths are generated, the antithetic-variates estimator averages over  $2N$  estimates.

## Variance Reduction: Antithetic Variates (continued)

- Consider process  $dX = a_t dt + b_t \sqrt{dt} \xi$ .
- Let  $g$  be a function of  $n$  samples  $X_1, X_2, \dots, X_n$  on the sample path.
- We are interested in  $E[g(X_1, X_2, \dots, X_n)]$ .
- Suppose one simulation run has realizations  $\xi_1, \xi_2, \dots, \xi_n$  for the normally distributed fluctuation term  $\xi$ .
- This generates samples  $x_1, x_2, \dots, x_n$ .
- The estimate is then  $g(\mathbf{x})$ , where  $\mathbf{x} \equiv (x_1, x_2, \dots, x_n)$ .

## Variance Reduction: Antithetic Variates (concluded)

- The antithetic-variates method does not sample  $n$  more numbers from  $\xi$  for the second estimate  $g(\mathbf{x}')$ .
- Instead, generate the sample path  $\mathbf{x}' \equiv (x'_1, x'_2, \dots, x'_n)$  from  $-\xi_1, -\xi_2, \dots, -\xi_n$ .
- Compute  $g(\mathbf{x}')$ .
- Output  $(g(\mathbf{x}) + g(\mathbf{x}'))/2$ .
- Repeat the above steps for as many times as required by accuracy.

## Variance Reduction: Conditioning

- We are interested in estimating  $E[X]$ .
- Suppose here is a random variable  $Z$  such that  $E[X | Z = z]$  can be efficiently and precisely computed.
- $E[X] = E[E[X | Z]]$  by the law of iterated conditional expectations.
- Hence the random variable  $E[X | Z]$  is also an unbiased estimator of  $E[X]$ .

## Variance Reduction: Conditioning (concluded)

- As

$$\text{Var}[E[X | Z]] \leq \text{Var}[X],$$

$E[X | Z]$  has a smaller variance than observing  $X$  directly.

- First obtain a random observation  $z$  on  $Z$ .
- Then calculate  $E[X | Z = z]$  as our estimate.
  - There is no need to resort to simulation in computing  $E[X | Z = z]$ .
- The procedure can be repeated a few times to reduce the variance.

## Control Variates

- Use the analytic solution of a similar yet simpler problem to improve the solution.
- Suppose we want to estimate  $E[X]$  and there exists a random variable  $Y$  with a known mean  $\mu \equiv E[Y]$ .
- Then  $W \equiv X + \beta(Y - \mu)$  can serve as a “controlled” estimator of  $E[X]$  for any constant  $\beta$ .
  - However  $\beta$  is chosen,  $W$  remains an unbiased estimator of  $E[X]$  as

$$E[W] = E[X] + \beta E[Y - \mu] = E[X].$$

## Control Variates (continued)

- Note that

$$\text{Var}[W] = \text{Var}[X] + \beta^2 \text{Var}[Y] + 2\beta \text{Cov}[X, Y], \quad (95)$$

- Hence  $W$  is less variable than  $X$  if and only if

$$\beta^2 \text{Var}[Y] + 2\beta \text{Cov}[X, Y] < 0. \quad (96)$$

## Control Variates (concluded)

- The success of the scheme clearly depends on both  $\beta$  and the choice of  $Y$ .
  - For example, arithmetic average-rate options can be priced by choosing  $Y$  to be the otherwise identical geometric average-rate option's price and  $\beta = -1$ .
- This approach is much more effective than the antithetic-variates method.



## Choice of $Y$

- In general, the choice of  $Y$  is ad hoc,<sup>a</sup> and experiments must be performed to confirm the wisdom of the choice.
- Try to match calls with calls and puts with puts.<sup>b</sup>
- On many occasions,  $Y$  is a discretized version of the derivative that gives  $\mu$ .
  - Discretely monitored geometric average-rate option vs. the continuously monitored geometric average-rate option given by formulas (36) on p. 384.

---

<sup>a</sup>But see Dai (B82506025, R86526008, D8852600), Chiu (R94922072), and Lyuu (2015).

<sup>b</sup>Contributed by Ms. Teng, Huei-Wen (R91723054) on May 25, 2004.

## Optimal Choice of $\beta$

- For some choices, the discrepancy can be significant, such as the lookback option.<sup>a</sup>
- Equation (95) on p. 786 is minimized when

$$\beta = -\text{Cov}[X, Y] / \text{Var}[Y].$$

– It is called beta in the book.

- For this specific  $\beta$ ,

$$\text{Var}[W] = \text{Var}[X] - \frac{\text{Cov}[X, Y]^2}{\text{Var}[Y]} = (1 - \rho_{X,Y}^2) \text{Var}[X],$$

where  $\rho_{X,Y}$  is the correlation between  $X$  and  $Y$ .

---

<sup>a</sup>Contributed by Mr. Tsai, Hwai (R92723049) on May 12, 2004.

## Optimal Choice of $\beta$ (continued)

- Note that the variance can never be increased with the optimal choice.
- Furthermore, the stronger  $X$  and  $Y$  are correlated, the greater the reduction in variance.
- For example, if this correlation is nearly perfect ( $\pm 1$ ), we could control  $X$  almost exactly.

## Optimal Choice of $\beta$ (continued)

- Typically, neither  $\text{Var}[Y]$  nor  $\text{Cov}[X, Y]$  is known.
- Therefore, we cannot obtain the maximum reduction in variance.
- We can guess these values and hope that the resulting  $W$  does indeed have a smaller variance than  $X$ .
- A second possibility is to use the simulated data to estimate these quantities.
  - How to do it efficiently in terms of time and space?

## Optimal Choice of $\beta$ (concluded)

- Observe that  $-\beta$  has the same sign as the correlation between  $X$  and  $Y$ .
- Hence, if  $X$  and  $Y$  are positively correlated,  $\beta < 0$ , then  $X$  is adjusted downward whenever  $Y > \mu$  and upward otherwise.
- The opposite is true when  $X$  and  $Y$  are negatively correlated, in which case  $\beta > 0$ .
- Suppose a suboptimal  $\beta + \epsilon$  is used instead.
- The variance increases by only  $\epsilon^2 \text{Var}[Y]$ .<sup>a</sup>

---

<sup>a</sup>Han and Lai (2010).

## A Pitfall

- A potential pitfall is to sample  $X$  and  $Y$  independently.
- In this case,  $\text{Cov}[X, Y] = 0$ .
- Equation (95) on p. 786 becomes

$$\text{Var}[W] = \text{Var}[X] + \beta^2 \text{Var}[Y].$$

- So whatever  $Y$  is, the variance is *increased!*
- Lesson:  $X$  and  $Y$  must be correlated.

## Problems with the Monte Carlo Method

- The error bound is only probabilistic.
- The probabilistic error bound of  $\sqrt{N}$  does not benefit from regularity of the integrand function.
- The requirement that the points be independent random samples are wasteful because of clustering.
- In reality, pseudorandom numbers generated by completely deterministic means are used.
- Monte Carlo simulation exhibits a great sensitivity on the seed of the pseudorandom-number generator.

# *Matrix Computation*



To set up a philosophy against physics is rash;  
philosophers who have done so  
have always ended in disaster.  
— Bertrand Russell

## Definitions and Basic Results

- Let  $A \equiv [a_{ij}]_{1 \leq i \leq m, 1 \leq j \leq n}$ , or simply  $A \in \mathbf{R}^{m \times n}$ , denote an  $m \times n$  matrix.
- It can also be represented as  $[a_1, a_2, \dots, a_n]$  where  $a_i \in \mathbf{R}^m$  are vectors.
  - Vectors are column vectors unless stated otherwise.
- $A$  is a square matrix when  $m = n$ .
- The rank of a matrix is the largest number of linearly independent columns.

## Definitions and Basic Results (continued)

- A square matrix  $A$  is said to be symmetric if  $A^T = A$ .
- A real  $n \times n$  matrix

$$A \equiv [a_{ij}]_{i,j}$$

is diagonally dominant if  $|a_{ii}| > \sum_{j \neq i} |a_{ij}|$  for  $1 \leq i \leq n$ .

- Such matrices are nonsingular.
- The identity matrix is the square matrix

$$I \equiv \text{diag}[1, 1, \dots, 1].$$

## Definitions and Basic Results (concluded)

- A matrix has full column rank if its columns are linearly independent.
- A real symmetric matrix  $A$  is positive definite if

$$x^T Ax = \sum_{i,j} a_{ij} x_i x_j > 0$$

for any nonzero vector  $x$ .

- A matrix  $A$  is positive definite if and only if there exists a matrix  $W$  such that  $A = W^T W$  and  $W$  has full column rank.

## Cholesky Decomposition

- Positive definite matrices can be factored as

$$A = LL^T,$$

called the Cholesky decomposition.

- Above,  $L$  is a lower triangular matrix.

## Generation of Multivariate Distribution

- Let  $\mathbf{x} \equiv [x_1, x_2, \dots, x_n]^T$  be a vector random variable with a positive definite covariance matrix  $C$ .
- As usual, assume  $E[\mathbf{x}] = \mathbf{0}$ .
- This covariance structure can be matched by  $P\mathbf{y}$ .
  - $C = PP^T$  is the Cholesky decomposition of  $C$ .<sup>a</sup>
  - $\mathbf{y} \equiv [y_1, y_2, \dots, y_n]^T$  is a vector random variable with a covariance matrix equal to the identity matrix.

---

<sup>a</sup>What if  $C$  is not positive definite? See Lai (R93942114) and Lyuu (2007).

## Generation of Multivariate Normal Distribution

- Suppose we want to generate the multivariate normal distribution with a covariance matrix  $C = PP^T$ .
  - First, generate independent standard normal distributions  $y_1, y_2, \dots, y_n$ .
  - Then

$$P[y_1, y_2, \dots, y_n]^T$$

has the desired distribution.

- These steps can then be repeated.

## Multivariate Derivatives Pricing

- Generating the multivariate normal distribution is essential for the Monte Carlo pricing of multivariate derivatives (pp. 710ff).
- For example, the rainbow option on  $k$  assets has payoff

$$\max(\max(S_1, S_2, \dots, S_k) - X, 0)$$

at maturity.

- The closed-form formula is a multi-dimensional integral.<sup>a</sup>

---

<sup>a</sup>Johnson (1987); Chen (D95723006) and Lyuu (2009).



## Multivariate Derivatives Pricing (concluded)

- Suppose  $dS_j/S_j = r dt + \sigma_j dW_j$ ,  $1 \leq j \leq k$ , where  $C$  is the correlation matrix for  $dW_1, dW_2, \dots, dW_k$ .
- Let  $C = PP^T$ .
- Let  $\xi$  consist of  $k$  independent random variables from  $N(0, 1)$ .
- Let  $\xi' = P\xi$ .
- Similar to Eq. (94) on p. 752,

$$S_{i+1} = S_i e^{(r - \sigma_j^2/2) \Delta t + \sigma_j \sqrt{\Delta t} \xi'_j}, \quad 1 \leq j \leq k.$$

## Least-Squares Problems

- The least-squares (LS) problem is concerned with

$$\min_{x \in \mathbf{R}^n} \| Ax - b \|,$$

where  $A \in \mathbf{R}^{m \times n}$ ,  $b \in \mathbf{R}^m$ ,  $m \geq n$ .

- The LS problem is called regression analysis in statistics and is equivalent to minimizing the mean-square error.
- Often written as

$$Ax = b.$$

## Polynomial Regression

- In polynomial regression,  $x_0 + x_1x + \cdots + x_nx^n$  is used to fit the data  $\{(a_1, b_1), (a_2, b_2), \dots, (a_m, b_m)\}$ .
- This leads to the LS problem,

$$\begin{bmatrix} 1 & a_1 & a_1^2 & \cdots & a_1^n \\ 1 & a_2 & a_2^2 & \cdots & a_2^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & a_m & a_m^2 & \cdots & a_m^n \end{bmatrix} \begin{bmatrix} x_0 \\ x_1 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} .$$

- Consult the text for solutions.

## American Option Pricing by Simulation

- The continuation value of an American option is the conditional expectation of the payoff from keeping the option alive now.
- The option holder must compare the immediate exercise value and the continuation value.
- In standard Monte Carlo simulation, each path is treated independently of other paths.
- But the decision to exercise the option cannot be reached by looking at one path alone.

## The Least-Squares Monte Carlo Approach

- The continuation value can be estimated from the cross-sectional information in the simulation by using least squares.<sup>a</sup>
- The result is a function (of the state) for estimating the continuation values.
- Use the function to estimate the continuation value for each path to determine its cash flow.
- This is called the least-squares Monte Carlo (LSM) approach.

---

<sup>a</sup>Longstaff and Schwartz (2001).

## The Least-Squares Monte Carlo Approach (concluded)

- The LSM is provably convergent.<sup>a</sup>
- The LSM can be easily parallelized.<sup>b</sup>
  - Partition the paths into subproblems and perform LSM on each of them independently.
  - The speedup is close to linear (i.e., proportional to the number of CPUs).
- Surprisingly, accuracy is not affected.

---

<sup>a</sup>Clément, Lamberton, and Protter (2002); Stentoft (2004).

<sup>b</sup>Huang (B96902079, R00922018) (2013) and Chen (B97902046, R01922005) (2014); Chen (B97902046, R01922005), Huang (B96902079, R00922018) (2013) and Lyuu (2015).

## A Numerical Example


- Consider a 3-year American put on a non-dividend-paying stock.
- The put is exercisable at years 0, 1, 2, and 3.
- The strike price  $X = 105$ .
- The annualized riskless rate is  $r = 5\%$ .
- The current stock price is 101.
  - The annual discount factor hence equals 0.951229.
- We use only 8 price paths to illustrate the algorithm.

## A Numerical Example (continued)

### Stock price paths

Path	Year 0	Year 1	Year 2	Year 3
1	101	<b>97.6424</b>	<b>92.5815</b>	107.5178
2	101	<b>101.2103</b>	105.1763	<b>102.4524</b>
3	101	105.7802	<b>103.6010</b>	124.5115
4	101	<b>96.4411</b>	<b>98.7120</b>	108.3600
5	101	124.2345	<b>101.0564</b>	<b>104.5315</b>
6	101	<b>95.8375</b>	<b>93.7270</b>	<b>99.3788</b>
7	101	108.9554	<b>102.4177</b>	<b>100.9225</b>
8	101	<b>104.1475</b>	113.2516	115.0994






## A Numerical Example (continued)

- We use the basis functions  $1, x, x^2$ .
  - Other basis functions are possible.<sup>a</sup>
- The plot next page shows the final estimated optimal exercise strategy given by LSM.
- We now proceed to tackle our problem.
- The idea is to calculate the cash flow along each path, using information from *all* paths.

---

<sup>a</sup>Laguerre polynomials, Hermite polynomials, Legendre polynomials, Chebyshev polynomials, Gedenbauer polynomials, and Jacobi polynomials.



## A Numerical Example (continued)

Cash flows at year 3

Path	Year 0	Year 1	Year 2	Year 3
1	—	—	—	0
2	—	—	—	2.5476
3	—	—	—	0
4	—	—	—	0
5	—	—	—	0.4685
6	—	—	—	5.6212
7	—	—	—	4.0775
8	—	—	—	0

## A Numerical Example (continued)

- The cash flows at year 3 are the exercise value if the put is in the money.
- Only 4 paths are in the money: 2, 5, 6, 7.
- Some of the cash flows may not occur if the put is exercised earlier, which we will find out step by step.
- Incidentally, the European counterpart has a value of

$$0.951229^3 \times \frac{2.5476 + 0.4685 + 5.6212 + 4.0775}{8} = 1.3680.$$

## A Numerical Example (continued)

- We move on to year 2.
- For each state that is in the money at year 2, we must decide whether to exercise it.
- There are 6 paths for which the put is in the money: 1, 3, 4, 5, 6, 7.
- Only in-the-money paths will be used in the regression because they are where early exercise is relevant.
  - If there were none, we would move on to year 1.

## A Numerical Example (continued)

- Let  $x$  denote the stock prices at year 2 for those 6 paths.
- Let  $y$  denote the corresponding discounted future cash flows (at year 3) if the put is not exercised at year 2.

## A Numerical Example (continued)

Regression at year 2

Path	$x$	$y$
1	92.5815	$0 \times 0.951229$
2	—	—
3	103.6010	$0 \times 0.951229$
4	98.7120	$0 \times 0.951229$
5	101.0564	$0.4685 \times 0.951229$
6	93.7270	$5.6212 \times 0.951229$
7	102.4177	$4.0775 \times 0.951229$
8	—	—



## A Numerical Example (continued)

- We regress  $y$  on 1,  $x$ , and  $x^2$ .
- The result is

$$f(x) = 22.08 - 0.313114 \times x + 0.00106918 \times x^2.$$

- $f(x)$  estimates the continuation value conditional on the stock price at year 2.
- We next compare the immediate exercise value and the continuation value.

## A Numerical Example (continued)

Optimal early exercise decision at year 2

Path	Exercise	Continuation
1	12.4185	$f(92.5815) = 2.2558$
2	—	—
3	1.3990	$f(103.6010) = 1.1168$
4	6.2880	$f(98.7120) = 1.5901$
5	3.9436	$f(101.0564) = 1.3568$
6	11.2730	$f(93.7270) = 2.1253$
7	2.5823	$f(102.4177) = 0.3326$
8	—	—

## A Numerical Example (continued)

- Amazingly, the put should be exercised in all 6 paths: 1, 3, 4, 5, 6, 7.
- Now, any positive cash flow at year 3 should be set to zero or overridden for these paths as the put is exercised before year 3.
  - They are paths 5, 6, 7.
- The cash flows on p. 815 become the ones on next slide.

## A Numerical Example (continued)

Cash flows at years 2 & 3

Path	Year 0	Year 1	Year 2	Year 3
1	—	—	12.4185	0
2	—	—	0	2.5476
3	—	—	1.3990	0
4	—	—	6.2880	0
5	—	—	3.9436	0
6	—	—	11.2730	0
7	—	—	2.5823	0
8	—	—	0	0

## A Numerical Example (continued)

- We move on to year 1.
- For each state that is in the money at year 1, we must decide whether to exercise it.
- There are 5 paths for which the put is in the money: 1, 2, 4, 6, 8.
- Only in-the-money paths will be used in the regression because they are where early exercise is relevant.
  - If there were none, we would move on to year 0.

## A Numerical Example (continued)

- Let  $x$  denote the stock prices at year 1 for those 5 paths.
- Let  $y$  denote the corresponding discounted future cash flows if the put is not exercised at year 1.
- From p. 823, we have the following table.

## A Numerical Example (continued)

Regression at year 1

Path	$x$	$y$
1	97.6424	$12.4185 \times 0.951229$
2	101.2103	$2.5476 \times 0.951229^2$
3	—	—
4	96.4411	$6.2880 \times 0.951229$
5	—	—
6	95.8375	$11.2730 \times 0.951229$
7	—	—
8	104.1475	0

## A Numerical Example (continued)

- We regress  $y$  on 1,  $x$ , and  $x^2$ .
- The result is

$$f(x) = -420.964 + 9.78113 \times x - 0.0551567 \times x^2.$$

- $f(x)$  estimates the continuation value conditional on the stock price at year 1.
- We next compare the immediate exercise value and the continuation value.



## A Numerical Example (continued)

Optimal early exercise decision at year 1

Path	Exercise	Continuation
1	7.3576	$f(97.6424) = 8.2230$
2	3.7897	$f(101.2103) = 3.9882$
3	—	—
4	8.5589	$f(96.4411) = 9.3329$
5	—	—
6	9.1625	$f(95.8375) = 9.83042$
7	—	—
8	0.8525	$f(104.1475) = -0.551885$

## A Numerical Example (continued)

- The put should be exercised for 1 path only: 8.
  - Note that  $f(104.1475) < 0$ .
- Now, any positive future cash flow should be set to zero or overridden for this path.
  - But there is none.
- The cash flows on p. 823 become the ones on next slide.
- They also confirm the plot on p. 814.

## A Numerical Example (continued)

Cash flows at years 1, 2, & 3

Path	Year 0	Year 1	Year 2	Year 3
1	—	0	12.4185	0
2	—	0	0	2.5476
3	—	0	1.3990	0
4	—	0	6.2880	0
5	—	0	3.9436	0
6	—	0	11.2730	0
7	—	0	2.5823	0
8	—	0.8525	0	0

## A Numerical Example (continued)

- We move on to year 0.
- The continuation value is, from p 830,

$$\begin{aligned} & (12.4185 \times 0.951229^2 + 2.5476 \times 0.951229^3 \\ & + 1.3990 \times 0.951229^2 + 6.2880 \times 0.951229^2 \\ & + 3.9436 \times 0.951229^2 + 11.2730 \times 0.951229^2 \\ & + 2.5823 \times 0.951229^2 + 0.8525 \times 0.951229) / 8 \\ = & 4.66263. \end{aligned}$$

## A Numerical Example (concluded)

- As this is larger than the immediate exercise value of

$$105 - 101 = 4,$$

the put should not be exercised at year 0.

- Hence the put's value is estimated to be 4.66263.
- Compare this with the European put's value of 1.3680 (p. 816).

# *Time Series Analysis*

The historian is a prophet in reverse.  
— Friedrich von Schlegel (1772–1829)

## GARCH Option Pricing<sup>a</sup>

- Options can be priced when the underlying asset's return follows a GARCH process.
- Let  $S_t$  denote the asset price at date  $t$ .
- Let  $h_t^2$  be the *conditional* variance of the return over the period  $[t, t + 1]$  given the information at date  $t$ .
  - “One day” is merely a convenient term for any elapsed time  $\Delta t$ .

---

<sup>a</sup>ARCH (autoregressive conditional heteroskedastic) is due to Engle (1982), co-winner of the 2003 Nobel Prize in Economic Sciences. GARCH (generalized ARCH ) is due to Bollerslev (1986) and Taylor (1986). A Bloomberg quant said to me on Feb 29, 2008, that GARCH is seldom used in trading.



## GARCH Option Pricing (continued)

- Adopt the following risk-neutral process for the price dynamics:<sup>a</sup>

$$\ln \frac{S_{t+1}}{S_t} = r - \frac{h_t^2}{2} + h_t \epsilon_{t+1}, \quad (97)$$

where

$$h_{t+1}^2 = \beta_0 + \beta_1 h_t^2 + \beta_2 h_t^2 (\epsilon_{t+1} - c)^2, \quad (98)$$

$$\epsilon_{t+1} \sim N(0, 1) \text{ given information at date } t,$$

$$r = \text{daily riskless return,}$$

$$c \geq 0.$$

---

<sup>a</sup>Duan (1995).

## GARCH Option Pricing (continued)

- The five unknown parameters of the model are  $c$ ,  $h_0$ ,  $\beta_0$ ,  $\beta_1$ , and  $\beta_2$ .
- It is postulated that  $\beta_0, \beta_1, \beta_2 \geq 0$  to make the conditional variance positive.
- There are other inequalities to satisfy (see text).
- The above process is called the nonlinear asymmetric GARCH (or NGARCH) model.

## GARCH Option Pricing (continued)

- It captures the volatility clustering in asset returns first noted by Mandelbrot (1963).<sup>a</sup>
  - When  $c = 0$ , a large  $\epsilon_{t+1}$  results in a large  $h_{t+1}$ , which in turns tends to yield a large  $h_{t+2}$ , and so on.
- It also captures the negative correlation between the asset return and changes in its (conditional) volatility.<sup>b</sup>
  - For  $c > 0$ , a positive  $\epsilon_{t+1}$  (good news) tends to decrease  $h_{t+1}$ , whereas a negative  $\epsilon_{t+1}$  (bad news) tends to do the opposite.

---

<sup>a</sup>“... large changes tend to be followed by large changes—of either sign—and small changes tend to be followed by small changes ...”

<sup>b</sup>Noted by Black (1976): Volatility tends to rise in response to “bad news” and fall in response to “good news.”

## GARCH Option Pricing (concluded)

- With  $y_t \equiv \ln S_t$  denoting the logarithmic price, the model becomes

$$y_{t+1} = y_t + r - \frac{h_t^2}{2} + h_t \epsilon_{t+1}. \quad (99)$$

- The pair  $(y_t, h_t^2)$  completely describes the current state.
- The conditional mean and variance of  $y_{t+1}$  are clearly

$$E[y_{t+1} | y_t, h_t^2] = y_t + r - \frac{h_t^2}{2}, \quad (100)$$

$$\text{Var}[y_{t+1} | y_t, h_t^2] = h_t^2. \quad (101)$$

## GARCH Model: Inferences

- Suppose the parameters  $c$ ,  $h_0$ ,  $\beta_0$ ,  $\beta_1$ , and  $\beta_2$  are given.
- Then we can recover  $h_1, h_2, \dots, h_n$  and  $\epsilon_1, \epsilon_2, \dots, \epsilon_n$  from the prices

$$S_0, S_1, \dots, S_n$$

under the GARCH model (97) on p. 836.

- This property is useful in statistical inferences.

## The Ritchken-Trevor (RT) Algorithm<sup>a</sup>

- The GARCH model is a continuous-state model.
- To approximate it, we turn to trees with *discrete* states.
- Path dependence in GARCH makes the tree for asset prices explode exponentially (why?).
- We need to mitigate this combinatorial explosion.

---

<sup>a</sup>Ritchken and Trevor (1999).

## The Ritchken-Trevor Algorithm (continued)

- Partition a day into  $n$  periods.
- Three states follow each state  $(y_t, h_t^2)$  after a period.
- As the trinomial model combines, each state at date  $t$  is followed by  $2n + 1$  states at date  $t + 1$  (recall p. 646).
- These  $2n + 1$  values must approximate the distribution of  $(y_{t+1}, h_{t+1}^2)$ .
- So the conditional moments (100)–(101) at date  $t + 1$  on p. 839 must be matched by the trinomial model to guarantee convergence to the continuous-state model.


## The Ritchken-Trevor Algorithm (continued)

- It remains to pick the jump size and the three branching probabilities.
- The role of  $\sigma$  in the Black-Scholes option pricing model is played by  $h_t$  in the GARCH model.
- As a jump size proportional to  $\sigma/\sqrt{n}$  is picked in the BOPM, a comparable magnitude will be chosen here.
- Define  $\gamma \equiv h_0$ , though other multiples of  $h_0$  are possible, and

$$\gamma_n \equiv \frac{\gamma}{\sqrt{n}}.$$

- The jump size will be some integer multiple  $\eta$  of  $\gamma_n$ .
- We call  $\eta$  the jump parameter (p. 844).





The seven values on the right approximate the distribution of logarithmic price  $y_{t+1}$ .

## The Ritchken-Trevor Algorithm (continued)

- The middle branch does not change the underlying asset's price.
- The probabilities for the up, middle, and down branches are

$$p_u = \frac{h_t^2}{2\eta^2\gamma^2} + \frac{r - (h_t^2/2)}{2\eta\gamma\sqrt{n}}, \quad (102)$$

$$p_m = 1 - \frac{h_t^2}{\eta^2\gamma^2}, \quad (103)$$

$$p_d = \frac{h_t^2}{2\eta^2\gamma^2} - \frac{r - (h_t^2/2)}{2\eta\gamma\sqrt{n}}. \quad (104)$$

## The Ritchken-Trevor Algorithm (continued)


- It can be shown that:
  - The trinomial model takes on  $2n + 1$  values at date  $t + 1$  for  $y_{t+1}$ .
  - These values have a matching mean for  $y_{t+1}$ .
  - These values have an asymptotically matching variance for  $y_{t+1}$ .
- The central limit theorem guarantees convergence as  $n$  increases (if the probabilities are valid).

## The Ritchken-Trevor Algorithm (continued)

- We can dispense with the intermediate nodes *between* dates to create a  $(2n + 1)$ -nomial tree (p. 848).
- The resulting model is multinomial with  $2n + 1$  branches from any state  $(y_t, h_t^2)$ .
- There are two reasons behind this manipulation.
  - Interdate nodes are created merely to approximate the continuous-state model after one day.
  - Keeping the interdate nodes results in a tree that can be  $n$  times larger.<sup>a</sup>

---

<sup>a</sup>Contrast it with the case on p. 366.



This heptanomial tree is the outcome of the trinomial tree on p. 844 after its intermediate nodes are removed.

## The Ritchken-Trevor Algorithm (continued)

- A node with logarithmic price  $y_t + \ell\eta\gamma_n$  at date  $t + 1$  follows the current node at date  $t$  with price  $y_t$ , where

$$-n \leq \ell \leq n.$$

- To reach that price in  $n$  periods, the number of up moves must exceed that of down moves by exactly  $\ell$ .
- The probability that this happens is

$$P(\ell) \equiv \sum_{j_u, j_m, j_d} \frac{n!}{j_u! j_m! j_d!} p_u^{j_u} p_m^{j_m} p_d^{j_d},$$

with  $j_u, j_m, j_d \geq 0$ ,  $n = j_u + j_m + j_d$ , and  $\ell = j_u - j_d$ .

## The Ritchken-Trevor Algorithm (continued)

- A particularly simple way to calculate the  $P(\ell)$ s starts by noting that

$$(p_u x + p_m + p_d x^{-1})^n = \sum_{\ell=-n}^n P(\ell) x^\ell. \quad (105)$$

- Convince yourself that this trick does the “accounting” correctly.
- So we expand  $(p_u x + p_m + p_d x^{-1})^n$  and retrieve the probabilities by reading off the coefficients.
- It can be computed in  $O(n^2)$  time, if not shorter.

## The Ritchken-Trevor Algorithm (continued)

- The updating rule (98) on p. 836 must be modified to account for the adoption of the discrete-state model.
- The logarithmic price  $y_t + \ell\eta\gamma_n$  at date  $t + 1$  following state  $(y_t, h_t^2)$  is associated with this variance:

$$h_{t+1}^2 = \beta_0 + \beta_1 h_t^2 + \beta_2 h_t^2 (\epsilon'_{t+1} - c)^2, \quad (106)$$

– Above,

$$\epsilon'_{t+1} = \frac{\ell\eta\gamma_n - (r - h_t^2/2)}{h_t}, \quad \ell = 0, \pm 1, \pm 2, \dots, \pm n,$$

is a discrete random variable with  $2n + 1$  values.



## The Ritchken-Trevor Algorithm (continued)

- Different conditional variances  $h_t^2$  may require different  $\eta$  so that the probabilities calculated by Eqs. (102)–(104) on p. 845 lie between 0 and 1.
- This implies varying jump sizes.
- The necessary requirement  $p_m \geq 0$  implies  $\eta \geq h_t/\gamma$ .
- Hence we try

$$\eta = \lceil h_t/\gamma \rceil, \lceil h_t/\gamma \rceil + 1, \lceil h_t/\gamma \rceil + 2, \dots$$

until valid probabilities are obtained or until their nonexistence is confirmed.

## The Ritchken-Trevor Algorithm (continued)


- The sufficient and necessary condition for valid probabilities to exist is<sup>a</sup>

$$\frac{|r - (h_t^2/2)|}{2\eta\gamma\sqrt{n}} \leq \frac{h_t^2}{2\eta^2\gamma^2} \leq \min\left(1 - \frac{|r - (h_t^2/2)|}{2\eta\gamma\sqrt{n}}, \frac{1}{2}\right).$$

- Obviously, the magnitude of  $\eta$  tends to grow with  $h_t$ .
- The plot on p. 854 uses  $n = 1$  to illustrate our points for a 3-day model.
- For example, node (1, 1) of date 1 and node (2, 3) of date 2 pick  $\eta = 2$ .

---

<sup>a</sup>Wu (R90723065) (2003); Lyuu and Wu (R90723065) (2003, 2005).



## The Ritchken-Trevor Algorithm (continued)

- The topology of the tree is not a standard combining multinomial tree.
- For example, a few nodes on p. 854 such as nodes  $(2, 0)$  and  $(2, -1)$  have *multiple* jump sizes.
- The reason is the path dependence of the model.
  - Two paths can reach node  $(2, 0)$  from the root node, each with a different variance for the node.
  - One of the variances results in  $\eta = 1$ , whereas the other results in  $\eta = 2$ .

## The Ritchken-Trevor Algorithm (concluded)

- The number of possible values of  $h_t^2$  at a node can be exponential.
  - Because each path brings a different variance  $h_t^2$ .
- To address this problem, we record only the maximum and minimum  $h_t^2$  at each node.<sup>a</sup>
- Therefore, each node on the tree contains only two states  $(y_t, h_{\max}^2)$  and  $(y_t, h_{\min}^2)$ .
- Each of  $(y_t, h_{\max}^2)$  and  $(y_t, h_{\min}^2)$  carries its own  $\eta$  and set of  $2n + 1$  branching probabilities.

---

<sup>a</sup>Cakici and Topyan (2000). But see p. 891 for a potential problem.

## Negative Aspects of the Ritchken-Trevor Algorithm<sup>a</sup>

- A small  $n$  may yield inaccurate option prices.
- But the tree will grow exponentially if  $n$  is large enough.
  - Specifically,  $n > (1 - \beta_1)/\beta_2$  when  $r = c = 0$ .
- A large  $n$  has another serious problem: The tree cannot grow beyond a certain date.
- Thus the choice of  $n$  may be quite limited in practice.
- The RT algorithm can be modified to be free of shortened maturity and exponential complexity.<sup>b</sup>

---

<sup>a</sup>Lyu and Wu (R90723065) (2003, 2005).

<sup>b</sup>Its size is only  $O(n^2)$  if  $n \leq (\sqrt{(1 - \beta_1)/\beta_2} - c)^2$ !