

Black-Scholes Differential Equation: An Alternative

- Perform the change of variable $V \triangleq \ln S$.
- The option value becomes $U(V, t) \triangleq C(e^V, t)$.
- Furthermore,

$$\begin{aligned}\frac{\partial C}{\partial t} &= \frac{\partial U}{\partial t}, \\ \frac{\partial C}{\partial S} &= \frac{1}{S} \frac{\partial U}{\partial V},\end{aligned}\tag{97}$$

$$\frac{\partial^2 C}{\partial S^2} = \frac{1}{S^2} \frac{\partial^2 U}{\partial V^2} - \frac{1}{S^2} \frac{\partial U}{\partial V}.\tag{98}$$

Black-Scholes Differential Equation: An Alternative (concluded)

- Equations (97) and (98) are alternative ways to calculate delta and gamma.^a
- They are very useful for trees of *logarithmic* prices.
- The Black-Scholes differential equation (95) on p. 689 becomes

$$\frac{1}{2} \sigma^2 \frac{\partial^2 U}{\partial V^2} + \left(r - q - \frac{\sigma^2}{2} \right) \frac{\partial U}{\partial V} - rU + \frac{\partial U}{\partial t} = 0$$

subject to $U(V, T)$ being the payoff such as $\max(X - e^V, 0)$.

^aRecall Eqs. (53) on p. 369 and (54) on p. 371.

PDEs for Asian Options

- Add the new variable $A(t) \triangleq \int_0^t S(u) du$.
- Then the value V of the Asian option satisfies this two-dimensional PDE:^a

$$\frac{\partial V}{\partial t} + rS \frac{\partial V}{\partial S} + \frac{1}{2} \sigma^2 S^2 \frac{\partial^2 V}{\partial S^2} + S \frac{\partial V}{\partial A} = rV.$$

- The terminal conditions are

$$V(T, S, A) = \max\left(\frac{A}{T} - X, 0\right) \quad \text{for call,}$$

$$V(T, S, A) = \max\left(X - \frac{A}{T}, 0\right) \quad \text{for put.}$$

^aKemna & Vorst (1990).

PDEs for Asian Options (continued)

- The two-dimensional PDE produces algorithms similar to that on pp. 451ff.^a
- But one-dimensional PDEs are available for Asian options.^b
- For example, Večer (2001) derives the following PDE for Asian calls:

$$\frac{\partial u}{\partial t} + r \left(1 - \frac{t}{T} - z \right) \frac{\partial u}{\partial z} + \frac{\left(1 - \frac{t}{T} - z \right)^2 \sigma^2}{2} \frac{\partial^2 u}{\partial z^2} = 0$$

with the terminal condition $u(T, z) = \max(z, 0)$.

^aBarraquand & Pudet (1996).

^bRogers & Shi (1995); Večer (2001); Dubois & Lelièvre (2005).

PDEs for Asian Options (concluded)

- For Asian puts:

$$\frac{\partial u}{\partial t} + r \left(\frac{t}{T} - 1 - z \right) \frac{\partial u}{\partial z} + \frac{\left(\frac{t}{T} - 1 - z \right)^2 \sigma^2}{2} \frac{\partial^2 u}{\partial z^2} = 0$$

with the same terminal condition.

- One-dimensional PDEs yield highly efficient numerical algorithms.

Hedging

When Professors Scholes and Merton and I
invested in warrants,
Professor Merton lost the most money.
And I lost the least.
— Fischer Black (1938–1995)

Delta Hedge

- Recall the delta (hedge ratio) of a derivative f :

$$\Delta \triangleq \frac{\partial f}{\partial S}.$$

- Thus

$$\Delta f \approx \Delta \times \Delta S$$

for relatively small changes in the stock price, ΔS .

- A delta-neutral portfolio is hedged as it is immunized against small changes in the stock price.

Delta Hedge (concluded)

- A trading strategy that dynamically maintains a delta-neutral portfolio is called delta hedge.
 - Trading strategies can also be static (or constant).^a
- Delta changes with the stock price.
- A delta hedge needs to be rebalanced periodically in order to maintain delta neutrality.
- In the limit where the portfolio is adjusted continuously, “perfect” hedge is achieved and the strategy becomes “self-financing.”

^aRecall p. 498 for one in hedging the short forward contract with the underlying asset and loans.

Implementing Delta Hedge

- We want to hedge N *short* derivatives.
- Assume the stock pays no dividends.
- The delta-neutral portfolio maintains $N \times \Delta$ shares of stock plus B borrowed dollars such that

$$-N \times f + N \times \Delta \times S - B = 0.$$

- At next rebalancing point when the delta is Δ' , buy $N \times (\Delta' - \Delta)$ shares to maintain $N \times \Delta'$ shares.
- Delta hedge is the discrete-time analog of the continuous-time limit.
- It will rarely be self-financing however small Δt is.

Example

- A hedger is *short* 10,000 European calls.
- $S = 50$, $\sigma = 30\%$, and $r = 6\%$.
- This call's expiration is four weeks away, its strike price is \$50, and each call has a current value of $f = 1.76791$.
- As an option covers 100 shares of stock, $N = 1,000,000$.
- The trader adjusts the portfolio weekly.
- The calls are replicated well if the cumulative cost of trading *stock* is close to the call premium's FV.^a

^aThis takes the replication viewpoint: One starts with zero dollar.

Example (continued)

- As $\Delta = 0.538560$

$$N \times \Delta = 538,560$$

shares are purchased for a total cost of

$$538,560 \times 50 = 26,928,000$$

dollars to make the portfolio delta-neutral.

- The trader finances the purchase by borrowing

$$B = N \times \Delta \times S - N \times f = 25,160,090$$

dollars net.^a

^aThis takes the hedging viewpoint: One starts with the option premium. The two viewpoints are equivalent. See Exercise 16.3.2 of the text.

Example (continued)

- At 3 weeks to expiration, the stock price rises to \$51.
- The new call value is $f' = 2.10580$.
- So before rebalancing, the portfolio is worth

$$-N \times f' + 538,560 \times 51 - Be^{0.06/52} = 171,622. \quad (99)$$

- The delta hedge is not self-financing as \$171,622 can be withdrawn.
 - It does *not* replicate the calls perfectly.

Example (continued)

- The magnitude of the tracking error^a can be mitigated if adjustments are made more frequently.
- The tracking error over *one* rebalancing act is positive about 68% of the time.
- Its expected value is ~ 0 under the *risk-neutral* probability measure.^b
 - But the stock price should be sampled under the *real-world* probability measure.^c

^aThe variation in the net portfolio value.

^bBoyle & Emanuel (1980).

^cRecall Eq. (94) on p. 687 or see p. 715.

Example (continued)

- The tracking error at maturity is proportional to vega.^a
- In practice tracking errors will cease to decrease beyond a certain rebalancing frequency.
- With a higher delta $\Delta' = 0.640355$, the trader buys

$$N \times (\Delta' - \Delta) = 101,795$$

shares for \$5,191,545.

- The number of shares is increased to $N \times \Delta' = 640,355$.

^aKamal & Derman (1999).

Example (continued)

- The cumulative cost is^a

$$26,928,000 \times e^{0.06/52} + 5,191,545 = 32,150,634.$$

- The portfolio is again delta-neutral.

^aWe take the replication viewpoint again. Under the BOPM, the replicating strategy is self-financing and matches the payoff perfectly.

τ	S	Option value f	Delta Δ	Change in delta	No. shares bought $N \times (5)$	Cost of shares $(1) \times (6)$	Cumulative cost $FV(8') + (7)$
	(1)	(2)	(3)	(5)	(6)	(7)	(8)
4	50	1.7679	0.53856	—	538,560	26,928,000	26,928,000
3	51	2.1058	0.64036	0.10180	101,795	5,191,545	32,150,634
2	53	3.3509	0.85578	0.21542	215,425	11,417,525	43,605,277
1	52	2.2427	0.83983	-0.01595	-15,955	-829,660	42,825,960
0	54	4.0000	1.00000	0.16017	160,175	8,649,450	51,524,853

- We take the replication viewpoint.
- The total number of shares is 1,000,000 at expiration (trading takes place at expiration, too).

Example (continued)

- At expiration, the trader has 1,000,000 shares.
- Pay off \$50,000,000, the in-the-money calls' total strike price.
- The trader is left with an obligation of

$$51,524,853 - 50,000,000 = 1,524,853,$$

which represents the replication cost.

- So if we had started with the PV of \$1,524,853, we would have replicated 10,000 such calls in *this* scenario.

Example (concluded)

- The FV of the call premium equals

$$1,767,910 \times e^{0.06 \times 4/52} = 1,776,088.$$

- That means the net gain in this scenario is

$$1,776,088 - 1,524,853 = 251,235$$

if we are hedging 10,000 short European calls.

Tracking Error Revisited

- Define the dollar gamma as $S^2\Gamma$.^a
- The change in value of a delta-hedged *long* option position after a duration of Δt is proportional to the dollar gamma.
- It is about

$$(1/2)S^2\Gamma[(\Delta S/S)^2 - \sigma^2\Delta t].$$

- See the plots on p. 658 and p. 660.
- $(\Delta S/S)^2$ is called the daily realized variance.

^aSee the gamma plot on p. 359.

Tracking Error Revisited (continued)

- In our particular case,

$$S = 50, \Gamma = 0.0957074, \Delta S = 1, \sigma = 0.3, \Delta t = 1/52.$$

- The estimated tracking error is

$$-(1/2) \times 50^2 \times 0.0957074 \times \left[(1/50)^2 - (0.09/52) \right] = 159,205.$$

- It is very close to our earlier number of 171,622.^a
- Delta hedge is also called gamma scalping.^b

^aRecall Eq. (99) on p. 704.

^bBennett (2014).

Tracking Error Revisited (continued)

- Let the rebalancing times be t_1, t_2, \dots, t_n .
- Let $\Delta S_i = S_{i+1} - S_i$.
- The total tracking error at expiration is about

$$\sum_{i=0}^{n-1} e^{r(T-t_i)} \frac{S_i^2 \Gamma_i}{2} \left[\left(\frac{\Delta S_i}{S_i} \right)^2 - \sigma^2 \Delta t \right].$$

- The tracking error is clearly path dependent.
- Mathematically,^a

$$\sum_{i=0}^{n-1} \left(\frac{\Delta S_i}{S_i} \right)^2 \rightarrow \sigma^2 T.$$

^aProtter (2005).

Tracking Error Revisited (concluded)^a

- The tracking error^b ϵ_n over n rebalancing acts has about the same probability of being positive as being negative.
- Subject to certain regularity conditions, the root-mean-square tracking error $\sqrt{E[\epsilon_n^2]}$ is $O(1/\sqrt{n})$.^c
- The root-mean-square tracking error increases with σ at first and then decreases.

^aBertsimas, Kogan, & Lo (2000).

^bSuch as 251,235 on p. 710.

^cGrannan & Swindle (1996).

Which Probability Measure?^a

- The profit and loss (i.e., tracking error) of a hedging strategy should be calculated under the *real-world* probability measure.
- But the deltas and option prices should be calculated under the *risk-neutral* probability measure.
- If whenever we sample the next stock price, backward induction is performed for the delta, it will take a long time to obtain the distribution of the profit and loss.
- How to do it efficiently?

^aContributed by Mr. Chiu, Tzu-Hsuan (R08723061) on April 9, 2021.

Delta-Gamma Hedge

- Delta hedge is based on the first-order approximation to changes in the derivative price, Δf , due to changes in the stock price, ΔS .
- When ΔS is not small, the second-order term, gamma $\Gamma \triangleq \partial^2 f / \partial S^2$, helps.
- A delta-gamma hedge is a delta hedge that maintains zero portfolio gamma; it is gamma neutral.
- To meet this extra condition, one more security needs to be brought in.

Delta-Gamma Hedge (concluded)

- Suppose we want to hedge short calls as before.
- A hedging call f_2 is brought in.
- To set up a delta-gamma hedge, we solve

$$\begin{aligned} -N \times f + n_1 \times S + n_2 \times f_2 - B &= 0 && \text{(self-financing),} \\ -N \times \Delta + n_1 + n_2 \times \Delta_2 - 0 &= 0 && \text{(delta neutrality),} \\ -N \times \Gamma + 0 + n_2 \times \Gamma_2 - 0 &= 0 && \text{(gamma neutrality),} \end{aligned}$$

for n_1 , n_2 , and B .

- The gammas of the stock and bond are 0.
- See the numerical example on pp. 231–232 of the text.

Other Hedges

- If volatility changes, delta-gamma hedge may not work well.
- An enhancement is the delta-gamma-vega hedge, which also maintains vega zero portfolio vega.
- To accomplish this, still one more security has to be brought into the process.
- In practice, delta-vega hedge, which may not maintain gamma neutrality, performs better than delta hedge.

Trees

I love a tree more than a man.
— Ludwig van Beethoven (1770–1827)

All those holes and pebbles.
Who could count them?
— James Joyce, *Ulysses* (1922)

And though the holes were rather small,
they had to count them all.
— The Beatles, *A Day in the Life* (1967)


The Combinatorial Method

- The combinatorial method can often cut the running time by an order of magnitude.
- The basic paradigm is to count the number of admissible paths that lead from the root to any terminal node.
- We first used this method in the linear-time algorithm for standard European option pricing.^a
- We will now apply it to price barrier options.

^aRecall p. 290.


The Reflection Principle^a

- Imagine a particle at position $(0, -\mathbf{a})$ on the integral lattice that is to reach $(n, -\mathbf{b})$.
- Without loss of generality, assume $\mathbf{a} > 0$ and $\mathbf{b} \geq 0$.
- This particle's movement:



- How many paths touch the x axis?

^aAndré (1887).



The Reflection Principle (continued)

- For a path from $(0, -\mathbf{a})$ to $(n, -\mathbf{b})$ that touches the x axis, let J denote the first point this happens.
- Reflect the portion of the path from $(0, -\mathbf{a})$ to J .
- A path from $(0, \mathbf{a})$ to $(n, -\mathbf{b})$ is constructed.
- It also hits the x axis at J for the first time.
- The one-to-one mapping shows the number of paths from $(0, -\mathbf{a})$ to $(n, -\mathbf{b})$ that touch the x axis equals the number of paths from $(0, \mathbf{a})$ to $(n, -\mathbf{b})$.

The Reflection Principle (concluded)

- A path of this kind has $(n + \mathbf{b} + \mathbf{a})/2$ down moves and $(n - \mathbf{b} - \mathbf{a})/2$ up moves.^a
- Hence there are

$$\binom{n}{\frac{n+\mathbf{a}+\mathbf{b}}{2}} = \binom{n}{\frac{n-\mathbf{a}-\mathbf{b}}{2}} \quad (100)$$

such paths for *even* $n + \mathbf{a} + \mathbf{b}$.

– Convention: $\binom{n}{k} = 0$ for $k < 0$ or $k > n$.

^aVerify it!

Pricing Barrier Options (Lyu, 1998)

- Focus on the down-and-in call with barrier $H < X$.
- So $H < S$.
- Define

$$a \triangleq \left\lceil \frac{\ln(X/(Sd^n))}{\ln(u/d)} \right\rceil = \left\lceil \frac{\ln(X/S)}{2\sigma\sqrt{\Delta t}} + \frac{n}{2} \right\rceil,$$
$$h \triangleq \left\lfloor \frac{\ln(H/(Sd^n))}{\ln(u/d)} \right\rfloor = \left\lfloor \frac{\ln(H/S)}{2\sigma\sqrt{\Delta t}} + \frac{n}{2} \right\rfloor.$$


- a is such that $\tilde{X} \triangleq Su^a d^{n-a}$ is the *terminal* price that is closest to X from above.
- h is such that $\tilde{H} \triangleq Su^h d^{n-h}$ is the *terminal* price that is closest to H from below.^a

^aSo we *underestimate* the option price.

Pricing Barrier Options (continued)

- The true barrier is replaced by the effective barrier \tilde{H} in the binomial model.
- A process with n moves hence ends up in the money if and only if the number of up moves is at least a .
- The price $Su^k d^{n-k}$ is at a distance of $2k$ from the lowest possible price Sd^n on the binomial tree as

$$Su^k d^{n-k} = Sd^{-k} d^{n-k} = Sd^{n-\boxed{2k}}. \quad (101)$$



Pricing Barrier Options (continued)

- A path from S to the terminal price $Su^j d^{n-j}$ has probability $p^j(1-p)^{n-j}$ of being taken.
- With reference to p. 728, the reflection principle (p. 723) can be applied with

$$a = n - 2h,$$

$$b = 2j - 2h,$$

in Eq. (100) on p. 725 by treating the \tilde{H} line as the x axis.

Pricing Barrier Options (continued)

- Therefore,

$$\binom{n}{\frac{n+(n-2h)+(2j-2h)}{2}} = \binom{n}{n-2h+j}$$

paths hit \tilde{H} in the process for $h \leq n/2$.

- The terminal price $Su^j d^{n-j}$ is reached by a path that hits the effective barrier with probability

$$\binom{n}{n-2h+j} p^j (1-p)^{n-j}, \quad j \leq 2h.$$

Pricing Barrier Options (concluded)

- The option value equals

$$\frac{\sum_{j=a}^{2h} \binom{n}{n-2h+j} p^j (1-p)^{n-j} (Su^j d^{n-j} - X)}{R^n}. \quad (102)$$

– $R \triangleq e^{r\tau/n}$ is the riskless return per period.

- It yields a linear-time algorithm.^a


^aLyuu (1998).

Convergence of BOPM

- Equation (102) results in the same sawtooth-like convergence shown on p. 412 (repeated on next page).
- The reasons are not hard to see.
- The effective barrier \tilde{H} rarely equals the true barrier H .

Convergence of BOPM (continued)

Down-and-in call value



Convergence of BOPM (continued)

- Convergence is actually good if we limit n to certain values—191, for example.
- These values make the true barrier coincide with or just above one of the stock price levels, that is,

$$H \approx Sd^j = Se^{-j\sigma\sqrt{\tau/n}}$$

for some integer j .

- The preferred n 's are thus

$$n = \left\lceil \frac{\tau}{(\ln(S/H)/(j\sigma))^2} \right\rceil, \quad j = 1, 2, 3, \dots$$

Convergence of BOPM (continued)

- There is only one minor technicality left.
- We picked the effective barrier to be one of the $n + 1$ possible *terminal* stock prices.
- However, the effective barrier above, Sd^j , corresponds to a terminal stock price only when $n - j$ is even.^a
- To close this gap, we decrement n by one, if necessary, to make $n - j$ an even number.

^aThis is because $j = n - 2k$ for some k by Eq. (101) on p. 727. Of course we could have adopted the more general form Sd^j ($-n \leq j \leq n$) for the effective barrier. It makes a good exercise.

Convergence of BOPM (concluded)


- The preferred n 's are now

$$n = \begin{cases} \ell, & \text{if } \ell - j \text{ is even,} \\ \ell - 1, & \text{otherwise,} \end{cases} \quad (103)$$

$j = 1, 2, 3, \dots$, where

$$\ell \triangleq \left\lfloor \frac{\tau}{(\ln(S/H)/(j\sigma))^2} \right\rfloor.$$

- Evaluate pricing formula (102) on p. 731 only with the n above.



Practical Implications^a

- This binomial model is $O(1/\sqrt{n})$ convergent in general but $O(1/n)$ convergent when the barrier is matched.^b
- Now that barrier options can be efficiently priced, we can afford to pick very large n (see next page).
- This has profound consequences.^c

^aLyu (1998).

^bJ. Lin (R95221010) (2008); J. Lin (R95221010) & Palmer (2013).

^cSee pp. 753ff.

n	Combinatorial method	
	Value	Time (milliseconds)
21	5.507548	0.30
84	5.597597	0.90
191	5.635415	2.00
342	5.655812	3.60
533	5.652253	5.60
768	5.654609	8.00
1047	5.658622	11.10
1368	5.659711	15.00
1731	5.659416	19.40
2138	5.660511	24.70
2587	5.660592	30.20
3078	5.660099	36.70
3613	5.660498	43.70
4190	5.660388	44.10
4809	5.659955	51.60
5472	5.660122	68.70
6177	5.659981	76.70
6926	5.660263	86.90
7717	5.660272	97.20

Practical Implications (concluded)

- Pricing is prohibitively time consuming when $S \approx H$ because

$$n \sim 1/\ln^2(S/H)$$

by formula (103) on 736.

- This is called the barrier-too-close problem.
- This observation is indeed true of standard quadratic-time binomial tree algorithms.
- But it no longer applies to linear-time algorithms (see next page).

Barrier at 95.0			Barrier at 99.5			Barrier at 99.9		
<i>n</i>	Value	Time	<i>n</i>	Value	Time	<i>n</i>	Value	Time
	.							
	:		795	7.47761	8	19979	8.11304	253
2743	2.56095	31.1	3184	7.47626	38	79920	8.11297	1013
3040	2.56065	35.5	7163	7.47682	88	179819	8.11300	2200
3351	2.56098	40.1	12736	7.47661	166	319680	8.11299	4100
3678	2.56055	43.8	19899	7.47676	253	499499	8.11299	6300
4021	2.56152	48.1	28656	7.47667	368	719280	8.11299	8500
True	2.5615			7.4767			8.1130	

(All times in milliseconds.)


Trinomial Tree

- Set up a trinomial approximation to the geometric Brownian motion^a

$$\frac{dS}{S} = r dt + \sigma dW.$$

- The three stock prices at time Δt are S , Su , and Sd , where $ud = 1$.
- Let the mean and variance of the stock price be SM and S^2V , respectively.

^aParkinson (1977); Boyle (1986).



Trinomial Tree (continued)

- By Eqs. (29) on p. 182,

$$M \triangleq e^{r\Delta t},$$
$$V \triangleq M^2(e^{\sigma^2\Delta t} - 1).$$

- Impose the matching of mean and that of variance:

$$1 = p_u + p_m + p_d,$$

$$SM = [p_u u + p_m + (p_d/u)] S,$$

$$S^2 V = p_u (Su - SM)^2 + p_m (S - SM)^2 + p_d (Sd - SM)^2.$$

Trinomial Tree (continued)

- Use linear algebra to verify that

$$p_u = \frac{u(V + M^2 - M) - (M - 1)}{(u - 1)(u^2 - 1)},$$
$$p_d = \frac{u^2(V + M^2 - M) - u^3(M - 1)}{(u - 1)(u^2 - 1)}.$$

- We must also make sure the probabilities lie between 0 and 1.

Trinomial Tree (concluded)

- There are countless variations.
- But all converge to the Black-Scholes option pricing model.^a
- Like the binomial model,^b the trinomial model has a linear-time algorithm for European options.^c

^aMadan, Milne, & Shefrin (1989).

^bRecall p. 290 and p. 731.

^cT. Chen (R94922003) (2007); J. Wang (R85526003), C. Wang (F95922018), T. Dai (B82506025, R86526008, D8852600), T. Chen (R94922003), L. Liu, & Zhou (2022).

A Trinomial Tree

- Use $u = e^{\lambda\sigma\sqrt{\Delta t}}$, where $\lambda \geq 1$ is a tunable parameter.
- Then

$$p_u \rightarrow \frac{1}{2\lambda^2} + \frac{(r + \sigma^2)\sqrt{\Delta t}}{2\lambda\sigma},$$
$$p_d \rightarrow \frac{1}{2\lambda^2} - \frac{(r - 2\sigma^2)\sqrt{\Delta t}}{2\lambda\sigma}.$$

- A nice choice for λ is $\sqrt{\pi/2}$.^a

^aOmberg (1988).

Barrier Options Revisited

- BOPM introduces a specification error by replacing the barrier with a nonidentical effective barrier.
- The trinomial model solves the problem by adjusting λ so that the barrier is hit exactly.^a

- When

$$S e^{-h\lambda\sigma\sqrt{\Delta t}} = H,$$

it takes h down moves to go from S to H , if h is an integer.

- Then

$$h = \frac{\ln(S/H)}{\lambda\sigma\sqrt{\Delta t}}.$$

^aRitchken (1995).

Barrier Options Revisited (continued)

- This is easy to achieve by adjusting λ .
- Typically, we find the smallest $\lambda \geq 1$ such that h is an integer.^a
 - Such a λ may not exist for very small n 's.^b
- Toward that end, we find the *largest* integer $j \geq 1$ that satisfies $\frac{\ln(S/H)}{j\sigma\sqrt{\Delta t}} \geq 1$ to be the h .
- Then let

$$\lambda = \frac{\ln(S/H)}{h\sigma\sqrt{\Delta t}}.$$

^aWhy must $\lambda \geq 1$?

^bThis is not hard to check.

Barrier Options Revisited (continued)

- Alternatively, simply pick

$$h = \left\lceil \frac{\ln(S/H)}{\sigma\sqrt{\Delta t}} \right\rceil.$$

- Make sure $h \geq 1$.
- Then let

$$\lambda = \frac{\ln(S/H)}{h\sigma\sqrt{\Delta t}}.$$

Barrier Options Revisited (concluded)

- This done, one of the layers of the trinomial tree coincides with the barrier.
- The following probabilities may be used,


$$p_u = \frac{1}{2\lambda^2} + \frac{\mu' \sqrt{\Delta t}}{2\lambda\sigma},$$

$$p_m = 1 - \frac{1}{\lambda^2},$$

$$p_d = \frac{1}{2\lambda^2} - \frac{\mu' \sqrt{\Delta t}}{2\lambda\sigma}.$$

$$- \mu' \triangleq r - (\sigma^2/2).$$

Down-and-in call value



Algorithms Comparison^a

- So which algorithm is better, binomial or trinomial?
- Algorithms are often compared based on the n value at which they “converge.”
 - The one with the smallest n wins.
- So giraffes are faster than cheetahs because they take fewer strides to travel the same distance!
- Performance must be based on actual running times, not n .^b

^aLyyu (1998).

^bPatterson & Hennessy (1994).

Algorithms Comparison (continued)

- Pages 733 and 752 seem to show the trinomial model converges at a smaller n than BOPM.
- It is in this sense when people say trinomial models converge faster than binomial ones.
- But does it make the trinomial model better then?

Algorithms Comparison (concluded)

- The linear-time binomial tree algorithm actually performs better than the trinomial one.
- See the next page, expanded from p. 739.
- The barrier-too-close problem is also too hard for a quadratic-time trinomial tree algorithm.^a
 - See pp. 766ff for an alternative solution.

^aLyu (1998).

n	Combinatorial method		Trinomial tree algorithm	
	Value	Time	Value	Time
21	5.507548	0.30		
84	5.597597	0.90	5.634936	35.0
191	5.635415	2.00	5.655082	185.0
342	5.655812	3.60	5.658590	590.0
533	5.652253	5.60	5.659692	1440.0
768	5.654609	8.00	5.660137	3080.0
1047	5.658622	11.10	5.660338	5700.0
1368	5.659711	15.00	5.660432	9500.0
1731	5.659416	19.40	5.660474	15400.0
2138	5.660511	24.70	5.660491	23400.0
2587	5.660592	30.20	5.660493	34800.0
3078	5.660099	36.70	5.660488	48800.0
3613	5.660498	43.70	5.660478	67500.0
4190	5.660388	44.10	5.660466	92000.0
4809	5.659955	51.60	5.660454	130000.0
5472	5.660122	68.70		
6177	5.659981	76.70		


(All times in milliseconds.)

Double-Barrier Options

- Double-barrier options are barrier options with two barriers $L < H$.
 - They make up “less than 5% of the light exotic market.”^a
- Assume $L < S < H$.
- The binomial model produces oscillating option values (see plot on next page).^b

^aBennett (2014).

^bChao (R86526053) (1999); T. Dai (B82506025, R86526008, D8852600) & Lyuu (2005).



Double-Barrier Options (concluded)

- The combinatorial method yields a linear-time algorithm.^a
- This binomial model is $O(1/\sqrt{n})$ convergent in general.^b
- If the barriers L and H depend on time, we have moving-barrier options.^c

^aSee p. 241 of the textbook.

^bGobet (1999).


^cRogers & Zane (1998).

Double-Barrier Knock-Out Options

- We knew how to pick the λ so that one of the layers of the trinomial tree coincides with one barrier, say H .
- This choice, however, does not guarantee that the other barrier, L , is also hit.
- One way to handle this problem is to *lower* the layer of the tree just above L to coincide with L .^a
 - More general ways to make the trinomial model hit both barriers are available.^b

^aRitchken (1995); Hull (1999).

^bHsu (R7526001, D89922012) & Lyuu (2006). T. Dai (B82506025, R86526008, D8852600) & Lyuu (2006) combine binomial and trinomial trees to derive an $O(n)$ -time algorithm for double-barrier options (see pp. 766ff).



Double-Barrier Knock-Out Options (continued)

- The probabilities of the nodes on the layer above L must be adjusted.
- Let ℓ be the positive integer such that

$$Sd^{\ell+1} < L < Sd^{\ell}.$$

- Hence the layer of the tree just above L has price Sd^{ℓ} .^a

^aYou probably cannot do the same thing for binomial models (why?).
Thanks to a lively discussion on April 25, 2012.

Double-Barrier Knock-Out Options (concluded)

- Define $\gamma > 1$ as the number satisfying

$$L = Sd^{\ell-1}e^{-\gamma\lambda\sigma\sqrt{\Delta t}}.$$

- The prices between the barriers are (from low to high)


$$L, Sd^{\ell-1}, \dots, Sd^2, Sd, S, Su, Su^2, \dots, Su^{h-1}, Su^h = H.$$

- The probabilities for the nodes with price equal to $Sd^{\ell-1}$ are


$$p'_u = \frac{b + a\gamma}{1 + \gamma}, \quad p'_d = \frac{b - a}{\gamma + \gamma^2}, \quad \text{and} \quad p'_m = 1 - p'_u - p'_d,$$

where $a \triangleq \mu'\sqrt{\Delta t}/(\lambda\sigma)$ and $b \triangleq 1/\lambda^2$.

Convergence: Binomial vs. Trinomial



Ideas for Binomial Trees To Handle Two Barriers




The Binomial-Trinomial Tree

- Replace the first step of the binomial tree with a trinomial structure for convergence and efficiency.^a
- The resulting tree is called the binomial-trinomial tree.^b
- Suppose a *binomial* tree will be built with Δt as the duration of one period.
- Node X at time t needs to pick three nodes on the binomial tree at time $t + \Delta t'$ as its successor nodes.
 - Later, $\Delta t \leq \Delta t' < 2\Delta t$.

^aT. Dai (B82506025, R86526008, D8852600) & Lyuu (2006, 2008, 2010).

^bThe idea first emerged in a hotel in Muroran, Hokkaido, Japan, in May of 2005.

The Binomial-Trinomial Tree (continued)



The Binomial-Trinomial Tree (continued)

- These three nodes should guarantee:
 1. The mean and variance of the stock price are matched.
 2. The branching probabilities are between 0 and 1.
- Let S_t be the stock price at node X (of time t).
- Use $s(z)$ to denote the stock price at node z .

The Binomial-Trinomial Tree (continued)

- Recall that the expected value of the logarithmic return $\ln(S_{t+\Delta t'}/S_t)$ at time $t + \Delta t'$ equals^a

$$\mu \triangleq \left(r - \frac{\sigma^2}{2} \right) \Delta t'. \quad (104)$$

- Its variance equals

$$\text{Var} \triangleq \sigma^2 \Delta t'. \quad (105)$$

- Let node B be the node whose logarithmic return $\hat{\mu} \triangleq \ln(s(\text{B})/S_t)$ is closest to μ among all the nodes at time $t + \Delta t'$.

^aRecall p. 305.

The Binomial-Trinomial Tree (continued)

- The middle branch from node X will end at node B.
- The two nodes A and C, which bracket node B, are the destinations of the other two branches from node X.
- Recall that adjacent nodes on the binomial tree are spaced at $2\sigma\sqrt{\Delta t}$ apart.
- Review the illustration on p. 767.

The Binomial-Trinomial Tree (continued)

- The three branching probabilities from node X are obtained through matching the mean and variance of the logarithmic return $\ln(S_{t+\Delta t'}/S_t)$.
- Recall that $\hat{\mu} = \ln(s(B)/S_t)$ is the logarithmic return of the middle node B.
- Let α , β , and γ be the differences between μ and the three logarithmic returns

$$\ln(s(A)/S_t), \ln(s(B)/S_t), \ln(s(C)/S_t),$$

in that order.

The Binomial-Trinomial Tree (continued)

- In other words,

$$\alpha \triangleq \hat{\mu} + 2\sigma\sqrt{\Delta t} - \mu = \beta + 2\sigma\sqrt{\Delta t}, \quad (106)$$

$$\beta \triangleq \hat{\mu} - \mu, \quad (107)$$

$$\gamma \triangleq \hat{\mu} - 2\sigma\sqrt{\Delta t} - \mu = \beta - 2\sigma\sqrt{\Delta t}. \quad (108)$$

- The three branching probabilities p_u, p_m, p_d then satisfy

$$p_u\alpha + p_m\beta + p_d\gamma = 0, \quad (109)$$

$$p_u\alpha^2 + p_m\beta^2 + p_d\gamma^2 = \text{Var}, \quad (110)$$

$$p_u + p_m + p_d = 1. \quad (111)$$

The Binomial-Trinomial Tree (concluded)

- Equation (109) matches the mean (104) of the logarithmic return $\ln(S_{t+\Delta t'}/S_t)$ on p. 769.
- Equation (110) matches its variance (105) on p. 769.
- The three probabilities can be proved to lie between 0 and 1 by Cramer's rule.

Pricing Double-Barrier Options


- Consider a double-barrier option with two barriers L and H , where $L < S < H$.
- We need to make each barrier coincide with a layer of the binomial tree for better convergence.
- The idea is to choose a Δt such that

$$\frac{\ln(H/L)}{2\sigma\sqrt{\Delta t}} \quad (112)$$

is a positive integer.

- The distance between two adjacent nodes such as nodes Y and Z in the figure on p. 775 is $2\sigma\sqrt{\Delta t}$.

Pricing Double-Barrier Options (continued)



Pricing Double-Barrier Options (continued)

- Suppose that the goal is a tree with $\sim m$ periods.
- Suppose we pick $\Delta\tau \triangleq T/m$ for the length of each period.
- There is no guarantee that $\frac{\ln(H/L)}{2\sigma\sqrt{\Delta\tau}}$ is an integer.
- Pick the largest $\Delta t \leq \Delta\tau$ which makes $\frac{\ln(H/L)}{2\sigma\sqrt{\Delta t}}$ some integer κ .
- In other words, we select

$$\Delta t = \left(\frac{\ln(H/L)}{2\kappa\sigma} \right)^2,$$

where $\kappa = \left\lceil \frac{\ln(H/L)}{2\sigma\sqrt{\Delta\tau}} \right\rceil$.

Pricing Double-Barrier Options (continued)

- We now proceed to build the binomial-trinomial tree.
- Start with the binomial part.
- Lay out the nodes from the low barrier L upward.
- Automatically, a layer coincides with the high barrier H .
- It is unlikely that Δt divides T .
- The initial stock price is also unlikely to be on a layer of the binomial tree.^a

^aRecall p. 775.

Pricing Double-Barrier Options (continued)

- The binomial-trinomial structure can address this problem as follows.
- Between time 0 and time T , the binomial tree spans $\lfloor T/\Delta t \rfloor - 1$ periods.
- Let the first period have a duration equal to

$$\Delta t' \triangleq T - \left(\left\lfloor \frac{T}{\Delta t} \right\rfloor - 1 \right) \Delta t.$$

- Then these $\lfloor T/\Delta t \rfloor$ periods span T years.
- It is easy to verify that $\Delta t \leq \Delta t' < 2\Delta t$.

Pricing Double-Barrier Options (continued)

- Start with the root node at time 0 and at a price with logarithmic return $\ln(S/S) = 0$.
- Find the three nodes on the binomial tree at time $\Delta t'$ as described earlier.
- Calculate the three branching probabilities to them.
- Grow the binomial tree from these three nodes until time T to obtain a binomial-trinomial tree with $\lfloor T/\Delta t \rfloor$ periods.
- Review the illustration on p. 775.

Pricing Double-Barrier Options (continued)

- Now the binomial-trinomial tree can be used to price double-barrier options by backward induction.
- That takes quadratic time.
- But a linear-time algorithm exists for double-barrier options on the *binomial* tree.^a
- Apply that algorithm to price the double-barrier option's prices at the three nodes at time $\Delta t'$.
 - That is, nodes A, B, and C on p. 775.
- Then calculate their expected discounted value at the root node.

^aSee p. 241 of the textbook; Chao (R86526053) (1999); T. Dai (B82506025, R86526008, D8852600) & Lyuu (2008).

Pricing Double-Barrier Options (continued)

- The overall running time is only linear!
- Binomial trees have troubles pricing barrier options.^a
- Even pit against the trinomial tree, the binomial-trinomial tree converges faster and smoother.^b
- In fact, the binomial-trinomial tree has an error of $O(1/n)$ for single-barrier options.^c
- Binomial trees with interpolation have an error of $O(1/n^{1-a})$ for any $0 < a < 1$ for double-barrier options.^d


^aSee p. 412, p. 758, and p. 764.

^bSee p. 782 and p. 783.

^cLyyu & Palmer (2010); Y. Lu (R06723032, D08922008) & Lyuu (2023).


^dAppolloni, Gaudenziy, & Zanette (2014).

Pricing Double-Barrier Options^a (continued)



^aGenerated by Mr. Lin, Ying-Hung (R01723029) on June 6, 2014.

Pricing Double-Barrier Options (concluded)



The thin line denotes the double-barrier option prices computed by the trinomial tree against the running time in seconds (such as point A). The thick line denotes those computed by the binomial-trinomial tree (such as point B).


The Barrier-Too-Close Problem (p. 740) Revisited

- Our idea solves it even if one barrier is very close to S .
 - It runs in linear time, unlike an earlier quadratic-time solution with trinomial trees (pp. 748ff).
 - Unlike an earlier solution using combinatorics (p. 731), now the choice of n is not that restricted.
- So it combines the strengths of binomial and trinomial trees.
- This holds for single-barrier options too.
- Here is how.

The Barrier-Too-Close Problem Revisited (continued)

- We can build the tree treating S as if it were a second barrier.
- So both H and S are matched.
- Alternatively, we can pick $\Delta\tau \triangleq T/m$ as our length of a period Δt without the subsequent adjustment.^a
- Then build the tree from the price H down.
- So H is matched.
- The initial price S is matched by the *trinomial* structure.

^aThere is no second barrier to match!



Plot of convergence of a down-and-out call to 5.9968 supplied by Mr. Lu, Yu-Ming (R06723032, D08922008) on February 26. 2021.

The Barrier-Too-Close Problem Revisited (concluded)

- The earlier trinomial tree is impractical as it needs a very large n when the barrier H is very close to S .^a
 - It needs at least one *up* move to connect S to H as its middle branch is flat.
 - But when $S \approx H$, that up move must take a very small step, necessitating a small Δt .
- Our trinomial structure's middle branch is *not* required to be flat.
- So S can be connected to H via the middle branch, and the need of a very large n disappears completely!

^aRecall the table on p. 741.